#### **Elementary Differential Equations 11th Edition Boyce Solutions Manual**

Full Download: https://alibabadownload.com/product/elementary-differential-equations-11th-edition-boyce-solutions-manual/



## Second-Order Linear Equations

1. Let  $y = e^{rt}$ , so that  $y' = r e^{rt}$  and  $y'' = r^2 e^{rt}$ . Direct substitution into the differential equation yields  $(r^2 + 2r - 3)e^{rt} = 0$ . Canceling the exponential, the characteristic equation is  $r^2 + 2r - 3 = 0$ . The roots of the equation are r = -3, 1. Hence the general solution is  $y = c_1 e^t + c_2 e^{-3t}$ .

2. Let  $y = e^{rt}$ . Substitution of the assumed solution results in the characteristic equation  $r^2 + 3r + 2 = 0$ . The roots of the equation are r = -2, -1. Hence the general solution is  $y = c_1 e^{-t} + c_2 e^{-2t}$ .

5. The characteristic equation is  $4r^2 - 9 = 0$ , with roots  $r = \pm 3/2$ . Therefore the general solution is  $y = c_1 e^{-3t/2} + c_2 e^{3t/2}$ .

6. The characteristic equation is  $r^2 - 2r - 2 = 0$ , with roots  $r = 1 \pm \sqrt{3}$ . Hence the general solution is  $y = c_1 e^{(1-\sqrt{3})t} + c_2 e^{(1+\sqrt{3})t}$ .

7. Substitution of the assumed solution  $y = e^{rt}$  results in the characteristic equation  $r^2 + r - 2 = 0$ . The roots of the equation are r = -2, 1. Hence the general solution is  $y = c_1 e^{-2t} + c_2 e^t$ . Its derivative is  $y' = -2c_1 e^{-2t} + c_2 e^t$ . Based on the first condition, y(0) = 1, we require that  $c_1 + c_2 = 1$ . In order to satisfy y'(0) = 1, we find that  $-2c_1 + c_2 = 1$ . Solving for the constants,  $c_1 = 0$  and  $c_2 = 1$ . Hence the specific solution is  $y(t) = e^t$ . It clearly increases without bound as  $t \to \infty$ .



9. The characteristic equation is  $r^2 + 3r = 0$ , with roots r = -3, 0. Therefore the general solution is  $y = c_1 + c_2 e^{-3t}$ , with derivative  $y' = -3 c_2 e^{-3t}$ . In order to satisfy the initial conditions, we find that  $c_1 + c_2 = -2$ , and  $-3 c_2 = 3$ . Hence the specific solution is  $y(t) = -1 - e^{-3t}$ . This converges to -1 as  $t \to \infty$ .



10. The characteristic equation is  $2r^2 + r - 4 = 0$ , with roots  $r = (-1 \pm \sqrt{33})/4$ . The general solution is  $y = c_1 e^{(-1-\sqrt{33})t/4} + c_2 e^{(-1+\sqrt{33})t/4}$ , with derivative

$$y' = \frac{-1 - \sqrt{33}}{4} c_1 e^{(-1 - \sqrt{33})t/4} + \frac{-1 + \sqrt{33}}{4} c_2 e^{(-1 + \sqrt{33})t/4}$$

In order to satisfy the initial conditions, we require that

$$c_1 + c_2 = 0$$
 and  $\frac{-1 - \sqrt{33}}{4}c_1 + \frac{-1 + \sqrt{33}}{4}c_2 = 1$ .

Solving for the coefficients,  $c_1=-2/\sqrt{33}$  and  $c_2=2/\sqrt{33}$  . The specific solution is

$$y(t) = -2 \left[ e^{(-1-\sqrt{33})t/4} - e^{(-1+\sqrt{33})t/4} \right] /\sqrt{33} .$$

It clearly increases without bound as  $t \to \infty$ .



12. The characteristic equation is  $4r^2 - 1 = 0$ , with roots  $r = \pm 1/2$ . Therefore the general solution is  $y = c_1 e^{-t/2} + c_2 e^{t/2}$ . Since the initial conditions are specified at t = -2, is more convenient to write  $y = d_1 e^{-(t+2)/2} + d_2 e^{(t+2)/2}$ . The derivative is given by  $y' = -\left[d_1 e^{-(t+2)/2}\right]/2 + \left[d_2 e^{(t+2)/2}\right]/2$ . In order to satisfy the initial conditions, we find that  $d_1 + d_2 = 1$ , and  $-d_1/2 + d_2/2 = -1$ . Solving for the coefficients,  $d_1 = 3/2$ , and  $d_2 = -1/2$ . The specific solution is

$$y(t) = \frac{3}{2}e^{-(t+2)/2} - \frac{1}{2}e^{(t+2)/2} = \frac{3}{2e}e^{-t/2} - \frac{e}{2}e^{t/2}$$

It clearly decreases without bound as  $t \to \infty$ .



15. The characteristic equation is  $2r^2 - 3r + 1 = 0$ , with roots r = 1/2, 1. Therefore the general solution is  $y = c_1 e^{t/2} + c_2 e^t$ , with derivative  $y' = c_1 e^{t/2}/2 + c_2 e^t$ . In order to satisfy the initial conditions, we require  $c_1 + c_2 = 2$  and  $c_1/2 + c_2 = 1/2$ . Solving for the coefficients,  $c_1 = 3$ , and  $c_2 = -1$ . The specific solution is  $y(t) = 3e^{t/2} - e^t$ . To find the stationary point, set  $y' = 3e^{t/2}/2 - e^t = 0$ . There is a unique solution, with  $t_1 = \ln(9/4)$ . The maximum value is then  $y(t_1) = 9/4$ . To find the *x*-intercept, solve the equation  $3e^{t/2} - e^t = 0$ . The solution is readily found to be  $t_2 = \ln 9 \approx 2.1972$ .

17. The characteristic equation is  $r^2 - (2\alpha - 1)r + \alpha(\alpha - 1) = 0$ . Examining the coefficients, the roots are  $r = \alpha$ ,  $\alpha - 1$ . Hence the general solution of the differential equation is  $y(t) = c_1 e^{\alpha t} + c_2 e^{(\alpha - 1)t}$ . Assuming  $\alpha \in \mathbb{R}$ , all solutions will tend to zero as long as  $\alpha < 0$ . On the other hand, all solutions will become unbounded as long as  $\alpha - 1 > 0$ , that is,  $\alpha > 1$ .

19.(a) The characteristic roots are r = -3, -2. The solution of the initial value problem is  $y(t) = (6 + \beta)e^{-2t} - (4 + \beta)e^{-3t}$ .

(b) The maximum point has coordinates  $t_0 = \ln [(3(4+\beta))/(2(6+\beta))], y_0 = 4(6+\beta)^3/(27(4+\beta)^2)$ .

- (c)  $y_0 = 4(6+\beta)^3/(27(4+\beta)^2) \ge 4$ , as long as  $\beta \ge 6+6\sqrt{3}$ .
- (d)  $\lim_{\beta \to \infty} t_0 = \ln(3/2)$ ,  $\lim_{\beta \to \infty} y_0 = \infty$ .

20.(a) Assuming that y is a constant, the differential equation reduces to cy = d. Hence the only equilibrium solution is y = d/c.

(b) Setting y = Y + d/c, substitution into the differential equation results in the equation aY'' + bY' + c(Y + d/c) = d. The equation satisfied by Y is aY'' + bY' + cY = 0.

1.

$$W(e^{2t}, e^{-3t/2}) = \begin{vmatrix} e^{2t} & e^{-3t/2} \\ 2e^{2t} & -\frac{3}{2}e^{-3t/2} \end{vmatrix} = -\frac{7}{2}e^{t/2}.$$

3.

$$W(e^{-2t}, t e^{-2t}) = \begin{vmatrix} e^{-2t} & t e^{-2t} \\ -2e^{-2t} & (1-2t)e^{-2t} \end{vmatrix} = e^{-4t}$$

4.

$$W(e^t \sin t, e^t \cos t) = \begin{vmatrix} e^t \sin t & e^t \cos t \\ e^t (\sin t + \cos t) & e^t (\cos t - \sin t) \end{vmatrix} = -e^{2t}.$$

5.

$$W(\cos^2\theta, 1 + \cos 2\theta) = \begin{vmatrix} \cos^2\theta & 1 + \cos 2\theta \\ -2\sin\theta\cos\theta & -2\sin 2\theta \end{vmatrix} = 0.$$

6. Write the equation as y'' + (3/t)y' = 1. p(t) = 3/t is continuous for all t > 0. Since  $t_0 > 0$ , the IVP has a unique solution for all t > 0.

7. Write the equation as y'' + (3/(t-4))y' + (4/t(t-4))y = 2/t(t-4). The coefficients are not continuous at t = 0 and t = 4. Since  $t_0 \in (0, 4)$ , the largest interval is 0 < t < 4.

8. The coefficient  $3 \ln |t|$  is discontinuous at t = 0. Since  $t_0 > 0$ , the largest interval of existence is  $0 < t < \infty$ .

10.  $y_1'' = 2$ . We see that  $t^2(2) - 2(t^2) = 0$ .  $y_2'' = 2t^{-3}$ , with  $t^2(y_2'') - 2(y_2) = 0$ . Let  $y_3 = c_1t^2 + c_2t^{-1}$ , then  $y_3'' = 2c_1 + 2c_2t^{-3}$ . It is evident that  $y_3$  is also a solution.

13. No. Substituting  $y = \sin(t^2)$  into the differential equation,

$$-4t^{2}\sin(t^{2}) + 2\cos(t^{2}) + 2t\cos(t^{2})p(t) + \sin(t^{2})q(t) = 0.$$

At t = 0, this equation becomes 2 = 0 (if we suppose that p(t) and q(t) are continuous), which is impossible.

14.  $W(e^{2t}, g(t)) = e^{2t}g'(t) - 2e^{2t}g(t) = 3e^{4t}$ . Dividing both sides by  $e^{2t}$ , we find that g must satisfy the ODE  $g' - 2g = 3e^{2t}$ . Hence  $g(t) = 3t e^{2t} + c e^{2t}$ .

15.  $W(f,g) = fg' - f'g = t \cos t - \sin t$ , and W(u,v) = -4fg' + 4f'g. Hence  $W(u,v) = -4t \cos t + 4 \sin t$ .

16. We compute

$$\begin{split} W(a_1y_1 + a_2y_2, b_1y_1 + b_2y_2) &= \begin{vmatrix} a_1y_1 + a_2y_2 & b_1y_1 + b_2y_2 \\ a_1y_1' + a_2y_2' & b_1y_1' + b_2y_2' \end{vmatrix} = \\ &= (a_1y_1 + a_2y_2)(b_1y_1' + b_2y_2') - (b_1y_1 + b_2y_2)(a_1y_1' + a_2y_2') = \\ &= a_1b_2(y_1y_2' - y_1'y_2) - a_2b_1(y_1y_2' - y_1'y_2) = (a_1b_2 - a_2b_1)W(y_1, y_2). \end{split}$$

This now readily shows that  $y_3$  and  $y_4$  form a fundamental set of solutions if and only if  $a_1b_2 - a_2b_1 \neq 0$ .

18. The general solution is  $y = c_1 e^{-3t} + c_2 e^{-t}$ .  $W(e^{-3t}, e^{-t}) = 2e^{-4t}$ , and hence the exponentials form a fundamental set of solutions. On the other hand, the fundamental solutions must also satisfy the conditions  $y_1(1) = 1$ ,  $y'_1(1) = 0$ ;  $y_2(1) = 0$ ,  $y'_2(1) = 1$ . For  $y_1$ , the initial conditions require  $c_1 + c_2 = e$ ,  $-3c_1 - c_2 = 0$ . The coefficients are  $c_1 = -e^3/2$ ,  $c_2 = 3e/2$ . For the solution  $y_2$ , the initial conditions require  $c_1 + c_2 = 0$ ,  $-3c_1 - c_2 = e$ . The coefficients are  $c_1 = -e^3/2$ ,  $c_2 = e/2$ . Hence the fundamental solutions are

$$y_1 = -\frac{1}{2}e^{-3(t-1)} + \frac{3}{2}e^{-(t-1)}$$
 and  $y_2 = -\frac{1}{2}e^{-3(t-1)} + \frac{1}{2}e^{-(t-1)}$ .

19. Yes.  $y_1'' = -4 \cos 2t$ ;  $y_2'' = -4 \sin 2t$ .  $W(\cos 2t, \sin 2t) = 2$ .

20. Clearly,  $y_1 = e^t$  is a solution.  $y'_2 = (1+t)e^t$ ,  $y''_2 = (2+t)e^t$ . Substitution into the ODE results in  $(2+t)e^t - 2(1+t)e^t + te^t = 0$ . Furthermore,  $W(e^t, te^t) = e^{2t}$ . Hence the solutions form a fundamental set of solutions.

24. Writing the equation in standard form, we find that  $P(t) = \sin t / \cos t$ . Hence the Wronskian is  $W(t) = c e^{-\int (\sin t / \cos t) dt} = c e^{\ln|\cos t|} = c \cos t$ , in which c is some constant.

25. Writing the equation in standard form, we find that  $P(x) = -2x/(1-x^2)$ . The Wronskian is  $W(x) = c e^{-\int -2x/(1-x^2) dx} = c e^{-\ln|1-x^2|} = c/(1-x^2)$ , in which c is some constant.

26. Rewrite the equation as p(t)y'' + p'(t)y' + q(t)y = 0. After writing the equation in standard form, we have P(t) = p'(t)/p(t). Hence the Wronskian is

$$W(t) = c e^{-\int p'(t)/p(t) dt} = c e^{-\ln p(t)} = c/p(t).$$

28. For the given differential equation, the Wronskian satisfies the first order differential equation W' + p(t)W = 0. Given that W is constant, it is necessary that  $p(t) \equiv 0$ .

32. P = 1, Q = x, R = 1. We have P'' - Q' + R = 0. The equation is exact. Note that (y')' + (xy)' = 0. Hence  $y' + xy = c_1$ . This equation is linear, with integrating factor  $\mu = e^{x^2/2}$ . Therefore the general solution is

$$y(x) = c_1 e^{-x^2/2} \int_{x_0}^x e^{u^2/2} du + c_2 e^{-x^2/2}.$$

34.  $P = x^2$ , Q = x, R = -1. We have P'' - Q' + R = 0. The equation is exact. Write the equation as  $(x^2y')' - (xy)' = 0$ . After integration, we conclude that  $x^2y' - xy = c$ . Divide both sides of the differential equation by  $x^2$ . The resulting equation is linear, with integrating factor  $\mu = 1/x$ . Hence  $(y/x)' = cx^{-3}$ . The solution is  $y(t) = c_1x^{-1} + c_2x$ .

36.  $P = x^2$ , Q = x,  $R = x^2 - \nu^2$ . Hence the coefficients are 2P' - Q = 3x and  $P'' - Q' + R = x^2 + 1 - \nu^2$ . The adjoint of the original differential equation is given by  $x^2\mu'' + 3x\mu' + (x^2 + 1 - \nu^2)\mu = 0$ .

37. P = 1, Q = 0, R = -x. Hence the coefficients are given by 2P' - Q = 0 and P'' - Q' + R = -x. Therefore the adjoint of the original equation is  $\mu'' - x \mu = 0$ .

### 3.3

- 1.  $e^{2-3i} = e^2 e^{-3i} = e^2 (\cos 3 i \sin 3).$
- 2.  $e^{i\pi} = \cos \pi + i \sin \pi = -1$ .
- 3.  $e^{2-(\pi/2)i} = e^2(\cos(\pi/2) i\sin(\pi/2)) = -e^2i$ .

6. The characteristic equation is  $r^2 - 2r + 6 = 0$ , with roots  $r = 1 \pm i\sqrt{5}$ . Hence the general solution is  $y = c_1 e^t \cos \sqrt{5}t + c_2 e^t \sin \sqrt{5}t$ .

7. The characteristic equation is  $r^2 + 2r + 2 = 0$ , with roots  $r = -1 \pm i$ . Hence the general solution is  $y = c_1 e^{-t} \cos t + c_2 e^{-t} \sin t$ .

9. The characteristic equation is  $r^2 + 2r + 1.25 = 0$ , with roots  $r = -1 \pm i/2$ . Hence the general solution is  $y = c_1 e^{-t} \cos(t/2) + c_2 e^{-t} \sin(t/2)$ .

11. The characteristic equation is  $r^2 + 4r + 6.25 = 0$ , with roots  $r = -2 \pm (3/2) i$ . Hence the general solution is  $y = c_1 e^{-2t} \cos(3t/2) + c_2 e^{-2t} \sin(3t/2)$ .

12. The characteristic equation is  $r^2 + 4 = 0$ , with roots  $r = \pm 2i$ . Hence the general solution is  $y = c_1 \cos 2t + c_2 \sin 2t$ . Now  $y' = -2c_1 \sin 2t + 2c_2 \cos 2t$ . Based on the first condition, y(0) = 0, we require that  $c_1 = 0$ . In order to satisfy the condition y'(0) = 1, we find that  $2c_2 = 1$ . The constants are  $c_1 = 0$  and  $c_2 = 1/2$ . Hence the specific solution is  $y(t) = \sin 2t/2$ . The solution is periodic.



13. The characteristic equation is  $r^2 - 2r + 5 = 0$ , with roots  $r = 1 \pm 2i$ . Hence the general solution is  $y = c_1 e^t \cos 2t + c_2 e^t \sin 2t$ . Based on the initial condition  $y(\pi/2) = 0$ , we require that  $c_1 = 0$ . It follows that  $y = c_2 e^t \sin 2t$ , and so the first derivative is  $y' = c_2 e^t \sin 2t + 2c_2 e^t \cos 2t$ . In order to satisfy the condition  $y'(\pi/2) = 2$ , we find that  $-2e^{\pi/2}c_2 = 2$ . Hence we have  $c_2 = -e^{-\pi/2}$ . Therefore the specific solution is  $y(t) = -e^{t-\pi/2} \sin 2t$ . The solution oscillates with an exponentially growing amplitude.



14. The characteristic equation is  $r^2 + 1 = 0$ , with roots  $r = \pm i$ . Hence the general solution is  $y = c_1 \cos t + c_2 \sin t$ . Its derivative is  $y' = -c_1 \sin t + c_2 \cos t$ . Based on the first condition,  $y(\pi/3) = 2$ , we require that  $c_1 + \sqrt{3}c_2 = 4$ . In order to satisfy the condition  $y'(\pi/3) = -4$ , we find that  $-\sqrt{3}c_1 + c_2 = -8$ . Solving

these for the constants,  $c_1 = 1 + 2\sqrt{3}$  and  $c_2 = \sqrt{3} - 2$ . Hence the specific solution is a steady oscillation, given by  $y(t) = (1 + 2\sqrt{3}) \cos t + (\sqrt{3} - 2) \sin t$ .



17.(a) The characteristic equation is  $5r^2 + 2r + 7 = 0$ , with roots  $r = -(1 \pm i\sqrt{34})/5$ . The solution is  $u = c_1 e^{-t/5} \cos \sqrt{34t}/5 + c_2 e^{-t/5} \sin \sqrt{34t}/5$ . Invoking the given initial conditions, we obtain the equations for the coefficients:  $c_1 = 2$ ,  $-2 + \sqrt{34} c_2 = 5$ . That is,  $c_1 = 2$ ,  $c_2 = 7/\sqrt{34}$ . Hence the specific solution is



(b) Based on the graph of u(t), T is in the interval 14 < t < 16. A numerical solution on that interval yields  $T \approx 14.5115$ .

19. Direct calculation gives the result. On the other hand, it can be shown that  $W(fg, fh) = f^2 W(g, h)$ . Hence  $W(e^{\lambda t} \cos \mu t, e^{\lambda t} \sin \mu t) = e^{2\lambda t} W(\cos \mu t, \sin \mu t) = e^{2\lambda t} [\cos \mu t (\sin \mu t)' - (\cos \mu t)' \sin \mu t] = \mu e^{2\lambda t}$ .

20.(a) Clearly,  $y_1$  and  $y_2$  are solutions. Also,  $W(\cos t, \sin t) = \cos^2 t + \sin^2 t = 1$ .

(b)  $y' = i e^{it}$ ,  $y'' = i^2 e^{it} = -e^{it}$ . Evidently, y is a solution and so  $y = c_1 y_1 + c_2 y_2$ .

(c) Setting t = 0,  $1 = c_1 \cos 0 + c_2 \sin 0$ , and  $c_1 = 1$ .

(d) Differentiating,  $i e^{it} = c_2 \cos t$ . Setting t = 0,  $i = c_2 \cos 0$  and hence  $c_2 = i$ . Therefore  $e^{it} = \cos t + i \sin t$ .

21. Euler's formula is  $e^{it} = \cos t + i \sin t$ . It follows that  $e^{-it} = \cos t - i \sin t$ . Adding these equation,  $e^{it} + e^{-it} = 2 \cos t$ . Subtracting the two equations results in  $e^{it} - e^{-it} = 2i \sin t$ .

22. Let 
$$r_1 = \lambda_1 + i\mu_1$$
, and  $r_2 = \lambda_2 + i\mu_2$ . Then  

$$e^{(r_1 + r_2)t} = e^{(\lambda_1 + \lambda_2)t + i(\mu_1 + \mu_2)t} = e^{(\lambda_1 + \lambda_2)t} \left[\cos(\mu_1 + \mu_2)t + i\sin(\mu_1 + \mu_2)t\right] =$$

$$= e^{(\lambda_1 + \lambda_2)t} \left[(\cos \mu_1 t + i\sin \mu_1 t)(\cos \mu_2 t + i\sin \mu_2 t)\right] =$$

$$= e^{\lambda_1 t} (\cos \mu_1 t + i\sin \mu_1 t) \cdot e^{\lambda_2 t} (\cos \mu_1 t + i\sin \mu_1 t) = e^{r_1 t} e^{r_2 t}.$$

Hence  $e^{(r_1+r_2)t} = e^{r_1t} e^{r_2t}$ .

23. Clearly,  $u' = \lambda e^{\lambda t} \cos \mu t - \mu e^{\lambda t} \sin \mu t = e^{\lambda t} (\lambda \cos \mu t - \mu \sin \mu t)$  and then  $u'' = \lambda e^{\lambda t} (\lambda \cos \mu t - \mu \sin \mu t) + e^{\lambda t} (-\lambda \mu \sin \mu t - \mu^2 \cos \mu t)$ . Plugging these into the differential equation, dividing by  $e^{\lambda t} \neq 0$  and arranging the sine and cosine terms we obtain that the identity to prove is

$$(a(\lambda^2 - \mu^2) + b\lambda + c)\cos\mu t + (-2\lambda\mu a - b\mu)\sin\mu t = 0.$$

We know that  $\lambda \pm i\mu$  solves the characteristic equation  $ar^2 + br + c = 0$ , so  $a(\lambda - i\mu)^2 + b(\lambda - i\mu) + c = a(\lambda^2 - \mu^2) + b\lambda + c + i(-2\lambda\mu a - \mu b) = 0$ . If this complex number is zero, then both the real and imaginary parts of it are zero, but those are the coefficients of  $\cos \mu t$  and  $\sin \mu t$  in the above identity, which proves that au'' + bu' + cu = 0. The solution for v is analogous.

26. The equation transforms into y'' + y = 0. The characteristic roots are  $r = \pm i$ . The solution is  $y = c_1 \cos(x) + c_2 \sin(x) = c_1 \cos(\ln t) + c_2 \sin(\ln t)$ .

28. The equation transforms into y'' - 5y' - 6y = 0. The characteristic roots are r = -1, 6. The solution is  $y = c_1 e^{-x} + c_2 e^{6x} = c_1 e^{-\ln t} + c_2 e^{6\ln t} = c_1/t + c_2 t^6$ .

29. The equation transforms into y'' - 5y' + 6y = 0. The characteristic roots are r = 2, 3. The solution is  $y = c_1 e^{2x} + c_2 e^{3x} = c_1 e^{2\ln t} + c_2 e^{3\ln t} = c_1 t^2 + c_2 t^3$ .

30. The equation transforms into y'' + 2y' - 3y = 0. The characteristic roots are r = 1, -3. The solution is  $y = c_1 e^x + c_2 e^{-3x} = c_1 e^{\ln t} + c_2 e^{-3\ln t} = c_1 t + c_2/t^3$ .

31. The equation transforms into y'' + 6y' + 10y = 0. The characteristic roots are  $r = -3 \pm i$ . The solution is

$$y = c_1 e^{-3x} \cos(x) + c_2 e^{-3x} \sin(x) = c_1 \frac{1}{t^3} \cos(\ln t) + c_2 \frac{1}{t^3} \sin(\ln t).$$

32.(a) By the chain rule, y'(x) = (dy/dx)x'. In general, dz/dt = (dz/dx)(dx/dt). Setting z = (dy/dt), we have

$$\frac{d^2y}{dt^2} = \frac{dz}{dx}\frac{dx}{dt} = \frac{d}{dx}\left[\frac{dy}{dx}\frac{dx}{dt}\right]\frac{dx}{dt} = \left[\frac{d^2y}{dx^2}\frac{dx}{dt}\right]\frac{dx}{dt} + \frac{dy}{dx}\frac{d}{dx}\left[\frac{dx}{dt}\right]\frac{dx}{dt}.$$

However,

$$\frac{d}{dx} \left[ \frac{dx}{dt} \right] \frac{dx}{dt} = \left[ \frac{d^2x}{dt^2} \right] \frac{dt}{dx} \cdot \frac{dx}{dt} = \frac{d^2x}{dt^2}$$

Hence

$$\frac{d^2y}{dt^2} = \frac{d^2y}{dx^2} \left[\frac{dx}{dt}\right]^2 + \frac{dy}{dx} \frac{d^2x}{dt^2}.$$

(b) Substituting the results in part (a) into the general differential equation, y'' + p(t)y' + q(t)y = 0, we find that

$$\frac{d^2y}{dx^2} \left[\frac{dx}{dt}\right]^2 + \frac{dy}{dx} \frac{d^2x}{dt^2} + p(t)\frac{dy}{dx} \frac{dx}{dt} + q(t)y = 0.$$

Collecting the terms,

$$\left[\frac{dx}{dt}\right]^2 \frac{d^2y}{dx^2} + \left[\frac{d^2x}{dt^2} + p(t)\frac{dx}{dt}\right]\frac{dy}{dx} + q(t)y = 0.$$

(c) Assuming  $(dx/dt)^2 = k q(t)$ , and q(t) > 0, we find that  $dx/dt = \sqrt{k q(t)}$ , which can be integrated. That is,  $x = u(t) = \int \sqrt{k q(t)} dt = \int \sqrt{q(t)} dt$ , since k = 1.

(d) Let k=1 . It follows that  $d^2x/dt^2 + p(t)dx/dt = du/dt + p(t)u(t) = q'/2\sqrt{q} + p\sqrt{q}$  . Hence

$$\left[\frac{d^2x}{dt^2} + p(t)\frac{dx}{dt}\right] / \left[\frac{dx}{dt}\right]^2 = \frac{q'(t) + 2p(t)q(t)}{2\left[q(t)\right]^{3/2}}.$$

As long as  $dx/dt \neq 0$ , the differential equation can be expressed as

$$\frac{d^2y}{dx^2} + \left\lfloor \frac{q'(t) + 2p(t)q(t)}{2[q(t)]^{3/2}} \right\rfloor \frac{dy}{dx} + y = 0.$$

(e) To find the analogue to the condition found in part d) for the case when q(t) < 0 we return to the conditions that make the coefficients on y, dy/dt and  $d^2y/dt^2$  proportional to each other. Since the coefficients on y and  $d^2y/dt^2$  are proportional,  $(dx/dt)^2 = \alpha q(t)$ , and we may take  $\alpha = -1$ . Thus  $dx/dt = (-q(t))^{1/2}$  and  $d^2y/dt^2 = (-q'/2)(-q)^{-1/2}$ . Since the coefficients on y and dy/dt are proportional, there is a constant  $\beta$  with

$$\beta q = \frac{d^2 y}{dt^2} + p(t)\frac{dx}{dt} = \frac{-q'}{2}(-q)^{-1/2} + p(-q)^{1/2} = \frac{-q'-2pq}{2(-q)^{1/2}}$$

and dividing each side of the equation by -q gives

$$-\beta = \frac{-q'-2pq}{2(-q)^{3/2}}, \text{ or } 2\beta = \frac{q'+2pq}{(-q)^{3/2}}$$

Thus the desired condition is that  $(q' + 2pq)/(-q)^{3/2}$  must be a constant.

34. Note that p(t) = 3t and  $q(t) = t^2$ . We have  $x = \int t \, dt = t^2/2$ . Furthermore,

$$\frac{q'(t) + 2p(t)q(t)}{2[q(t)]^{3/2}} = \frac{1+3t^2}{t^2}$$

The ratio is not constant, and therefore the equation cannot be transformed.

35. Note that p(t) = t - 1/t and  $q(t) = t^2$ . We have  $x = \int t \, dt = t^2/2$ . Furthermore,

$$\frac{q'(t) + 2p(t)q(t)}{2[q(t)]^{3/2}} = 1.$$

The ratio is constant, and therefore the equation can be transformed. From Problem 32, the transformed equation is

$$\frac{d^2y}{dx^2} + \frac{dy}{dx} + y = 0.$$

Based on the methods in this section, the characteristic equation is  $r^2 + r + 1 = 0$ , with roots  $r = (-1 \pm i\sqrt{3})/2$ . The general solution is  $y(x) = c_1 e^{-x/2} \cos \sqrt{3} x/2 + c_2 e^{-x/2} \sin \sqrt{3} x/2$ . Since  $x = t^2/2$ , the solution in the original variable t is

$$y(t) = e^{-t^2/4} \left[ c_1 \cos\left(\sqrt{3} t^2/4\right) + c_2 \sin\left(\sqrt{3} t^2/4\right) \right].$$

36. Note that p(t) = t and  $q(t) = -e^{-t^2} < 0$  for  $-\infty < t < \infty$ . To proceed we must confirm that  $(q' + 2pq)/(-q)^{3/2}$  is a constant:

$$\frac{q'+2pq}{(-q)^{3/2}} = \frac{2te^{-t^2}+2t(-e^{t^2})}{(e^{-t^2})^{3/2}} = 0$$

Thus the differential equation can be transformed into an equation with constant coefficients by letting  $x = u(t) = \int e^{-t^2/2} dt$ . Substituting x = u(t) in the differential equation found in part (b) of Problem 32 we obtain, after dividing by the coefficient of  $d^2y/dx^2$ , the differential equation  $(d^2y/dx^2) - y = 0$ . Hence the general solution of the original differential equation is  $y(t) = c_1 e^{x(t)} + c_2 e^{-x(t)}$ , where  $x(t) = \int e^{-t^2/2} dt$ .

2. The characteristic equation is  $9r^2 + 6r + 1 = 0$ , with the double root r = -1/3. The general solution is  $y(t) = c_1 e^{-t/3} + c_2 t e^{-t/3}$ .

3. The characteristic equation is  $4r^2 - 4r - 3 = 0$ , with roots r = -1/2, 3/2. The general solution is  $y(t) = c_1 e^{-t/2} + c_2 e^{3t/2}$ .

5. The characteristic equation is  $r^2 - 6r + 9 = 0$ , with the double root r = 3. The general solution is  $y(t) = c_1 e^{3t} + c_2 t e^{3t}$ .

3.4

6. The characteristic equation is  $4r^2 + 17r + 4 = 0$ , with roots r = -1/4, -4. The general solution is  $y(t) = c_1 e^{-t/4} + c_2 e^{-4t}$ .

7. The characteristic equation is  $16r^2 + 24r + 9 = 0$ , with double root r = -3/4. The general solution is  $y(t) = c_1 e^{-3t/4} + c_2 t e^{-3t/4}$ .

8. The characteristic equation is  $2r^2 + 2r + 1 = 0$ . We obtain the complex roots  $r = (-1 \pm i)/2$ . The general solution is  $y(t) = c_1 e^{-t/2} \cos(t/2) + c_2 e^{-t/2} \sin(t/2)$ .

9. The characteristic equation is  $9r^2 - 12r + 4 = 0$ , with the double root r = 2/3. The general solution is  $y(t) = c_1 e^{2t/3} + c_2 t e^{2t/3}$ . Invoking the first initial condition, it follows that  $c_1 = 2$ . Now  $y'(t) = (4/3 + c_2)e^{2t/3} + 2c_2 t e^{2t/3}/3$ . Invoking the second initial condition,  $4/3 + c_2 = -1$ , or  $c_2 = -7/3$ . Hence we obtain the solution  $y(t) = 2e^{2t/3} - (7/3)te^{2t/3}$ . Since the second term dominates for large t,  $y(t) \to -\infty$ .



12. The characteristic roots are  $r_1 = r_2 = 1/2$ . Hence the general solution is given by  $y(t) = c_1 e^{t/2} + c_2 t e^{t/2}$ . Invoking the initial conditions, we require that  $c_1 = 2$ , and that  $1 + c_2 = b$ . The specific solution is  $y(t) = 2e^{t/2} + (b-1)t e^{t/2}$ . Since the second term dominates, the long-term solution depends on the sign of the coefficient b-1. The critical value is b = 1.

15.(a) The characteristic equation is  $r^2 + 2ar + a^2 = (r+a)^2 = 0$ .

(b) With p(t) = 2a, Abel's Formula becomes  $W(y_1, y_2) = c e^{-\int 2a dt} = c e^{-2at}$ .

(c)  $y_1(t) = e^{-at}$  is a solution. From part (b), with c = 1,  $e^{-at} y_2'(t) + a e^{-at} y_2(t) = e^{-2at}$ , which can be written as  $(e^{at} y_2(t))' = 1$ , resulting in  $e^{at} y_2(t) = t$ .

17.(a) If the characteristic equation  $ar^2 + br + c$  has equal roots  $r_1$ , then  $ar_1^2 + br_1 + c = a(r - r_1)^2 = 0$ . Then clearly  $L[e^{rt}] = (ar^2 + br + c)e^{rt} = a(r - r_1)^2e^{rt}$ . This gives immediately that  $L[e^{r_1t}] = 0$ .

(b) Differentiating the identity in part (a) with respect to r we get  $(2ar + b)e^{rt} + (ar^2 + br + c)te^{rt} = 2a(r - r_1)e^{rt} + a(r - r_1)^2te^{rt}$ . Again, this gives  $L[te^{r_1t}] = 0$ .

18. Set  $y_2(t) = t^2 v(t)$ . Substitution into the differential equation results in

$$t^{2}(t^{2}v'' + 4tv' + 2v) - 4t(t^{2}v' + 2tv) + 6t^{2}v = 0.$$

After collecting terms, we end up with  $t^4v'' = 0$ . Hence  $v(t) = c_1 + c_2t$ , and thus  $y_2(t) = c_1t^2 + c_2t^3$ . Setting  $c_1 = 0$  and  $c_2 = 1$ , we obtain  $y_2(t) = t^3$ .

19. Set  $y_2(t) = t v(t)$ . Substitution into the differential equation results in

$$t^{2}(tv'' + 2v') + 2t(tv' + v) - 2tv = 0.$$

After collecting terms, we end up with  $t^3v'' + 4t^2v' = 0$ . This equation is linear in the variable w = v'. It follows that  $v'(t) = ct^{-4}$ , and  $v(t) = c_1t^{-3} + c_2$ . Thus  $y_2(t) = c_1t^{-2} + c_2t$ . Setting  $c_1 = 1$  and  $c_2 = 0$ , we obtain  $y_2(t) = t^{-2}$ .

23. Direct substitution verifies that  $y_1(t) = e^{-\delta x^2/2}$  is a solution of the differential equation. Now set  $y_2(x) = y_1(x) v(x)$ . Substitution of  $y_2$  into the equation results in  $v'' - \delta x v' = 0$ . This equation is linear in the variable w = v'. An integrating factor is  $\mu = e^{-\delta x^2/2}$ . Rewrite the equation as  $[e^{-\delta x^2/2} v']' = 0$ , from which it follows that  $v'(x) = c_1 e^{\delta x^2/2}$ . Integrating, we obtain

$$v(x) = c_1 \int_0^x e^{\delta u^2/2} du + v(0).$$

Hence

$$y_2(x) = c_1 e^{-\delta x^2/2} \int_0^x e^{\delta u^2/2} du + c_2 e^{-\delta x^2/2}.$$

Setting  $c_2 = 0$ , we obtain a second independent solution.

25. After writing the differential equation in standard form, we have p(t) = 3/t. Based on Abel's identity,  $W(y_1, y_2) = c_1 e^{-\int 3/t \, dt} = c_1 t^{-3}$ . As shown in Problem 24, two solutions of a second order linear equation satisfy  $(y_2/y_1)' = W(y_1, y_2)/y_1^2$ . In the given problem,  $y_1(t) = t^{-1}$ . Hence  $(t y_2)' = c_1 t^{-1}$ . Integrating both sides of the equation,  $y_2(t) = c_1 t^{-1} \ln t + c_2 t^{-1}$ . Setting  $c_1 = 1$  and  $c_2 = 0$  we obtain  $y_2(t) = t^{-1} \ln t$ .

27. Write the differential equation in standard form to find p(x) = 1/x. Based on Abel's identity,  $W(y_1, y_2) = c e^{-\int 1/x \, dx} = c x^{-1}$ . Two solutions of a second order linear differential equation satisfy  $(y_2/y_1)' = W(y_1, y_2)/y_1^2$ . In the given problem,  $y_1(x) = x^{-1/2} \sin x$ . Hence

$$\left(\frac{\sqrt{x}}{\sin x}\,y_2\right)' = c\,\frac{1}{\sin^2 x}\,.$$

Integrating both sides of the equation,  $y_2(x) = c_1 x^{-1/2} \cos x + c_2 x^{-1/2} \sin x$ . Setting  $c_1 = 1$  and  $c_2 = 0$ , we obtain  $y_2(x) = x^{-1/2} \cos x$ .

29.(a) The characteristic equation is  $ar^2+c=0\,.$  If  $a\,,c>0\,,$  then the roots are  $r=\pm\,i\sqrt{c/a}$  . The general solution is

$$y(t) = c_1 \cos \sqrt{\frac{c}{a}} t + c_2 \sin \sqrt{\frac{c}{a}} t$$
,

which is bounded.

(b) The characteristic equation is  $ar^2 + br = 0$ . The roots are r = 0, -b/a, and hence the general solution is  $y(t) = c_1 + c_2 e^{-bt/a}$ . Clearly,  $y(t) \to c_1$ . With the given initial conditions,  $c_1 = y_0 + (a/b)y'_0$ .

30. Note that  $2 \cos t \sin t = \sin 2t$ . Then  $1 - k \cos t \sin t = 1 - (k/2) \sin 2t$ . Now if 0 < k < 2, then  $(k/2) \sin 2t < |\sin 2t|$  and  $-(k/2) \sin 2t > -|\sin 2t|$ . Hence

$$1 - k \cos t \sin t = 1 - \frac{k}{2} \sin 2t > 1 - |\sin 2t| \ge 0.$$

31. The equation transforms into y'' - 4y' + 4y = 0. We obtain a double root r = 2. The solution is  $y = c_1 e^{2x} + c_2 x e^{2x} = c_1 e^{2 \ln t} + c_2 \ln t e^{2 \ln t} = c_1 t^2 + c_2 t^2 \ln t$ .

33. The equation transforms into y'' + 2y' + y = 0. We get a double root r = -1. The solution is  $y = c_1 e^{-x} + c_2 x e^{-x} = c_1 e^{-\ln t} + c_2 \ln t e^{-\ln t} = c_1 t^{-1} + c_2 t^{-1} \ln t$ .

34. The equation transforms into y'' - 3y' + 9y/4 = 0. We obtain the double root r = 3/2. The solution is  $y = c_1 e^{3x/2} + c_2 x e^{3x/2} = c_1 e^{3\ln t/2} + c_2 \ln t e^{3\ln t/2} = c_1 t^{3/2} + c_2 t^{3/2} \ln t$ .

# 3.5

2. The characteristic equation for the homogeneous problem is  $r^2 - r - 2 = 0$ , with roots r = -1, 2. Hence  $y_c(t) = c_1 e^{-t} + c_2 e^{2t}$ . Set  $Y = At^2 + Bt + C$ . Substitution into the given differential equation, and comparing the coefficients, results in the system of equations -2A = 4, -2A - 2B = -2 and 2A - B - 2C = 0. Hence  $Y = -2t^2 + 3t - 7/2$ . The general solution is  $y(t) = y_c(t) + Y$ .

3. The characteristic equation for the homogeneous problem is  $r^2 + r - 6 = 0$ , with roots r = -3, 2. Hence  $y_c(t) = c_1 e^{-3t} + c_2 e^{2t}$ . Set  $Y = Ae^{3t} + Be^{-2t}$ . Substitution into the given differential equation, and comparing the coefficients, results in the system of equations 6A = 12 and -4B = 12. Hence  $Y = 2e^{3t} - 3e^{-2t}$ . The general solution is  $y(t) = y_c(t) + Y$ .

4. The characteristic equation for the homogeneous problem is  $r^2 - 2r - 3 = 0$ , with roots r = -1, 3. Hence  $y_c(t) = c_1 e^{-t} + c_2 e^{3t}$ . Note that the assignment  $Y = Ate^{-t}$  is not sufficient to match the coefficients. Try  $Y = Ate^{-t} + Bt^2e^{-t}$ . Substitution into the differential equation, and comparing the coefficients, results in the system of equations -4A + 2B = 0 and -8B = -3. This implies that  $Y = (3/16)te^{-t} + (3/8)t^2e^{-t}$ . The general solution is  $y(t) = y_c(t) + Y$ .

8. The characteristic equation for the homogeneous problem is  $r^2 + \omega_0^2 = 0$ , with complex roots  $r = \pm \omega_0 i$ . Hence  $y_c(t) = c_1 \cos \omega_0 t + c_2 \sin \omega_0 t$ . Since  $\omega \neq \omega_0$ , set  $Y = A \cos \omega t + B \sin \omega t$ . Substitution into the ODE and comparing the coefficients results in the system of equations  $(\omega_0^2 - \omega^2)A = 1$  and  $(\omega_0^2 - \omega^2)B = 0$ .

Hence

$$Y = \frac{1}{\omega_0^2 - \omega^2} \cos \omega t \,.$$

The general solution is  $y(t) = y_c(t) + Y$ .

9. From Problem 8,  $y_c(t)$  is known. Since  $\cos \omega_0 t$  is a solution of the homogeneous problem, set  $Y = At \cos \omega_0 t + Bt \sin \omega_0 t$ . Substitution into the given ODE and comparing the coefficients results in A = 0 and  $B = 1/2\omega_0$ . Hence the general solution is  $y(t) = c_1 \cos \omega_0 t + c_2 \sin \omega_0 t + t \sin \omega_0 t/(2\omega_0)$ .

12. The characteristic equation for the homogeneous problem is  $r^2 + 4 = 0$ , with roots  $r = \pm 2i$ . Hence  $y_c(t) = c_1 \cos 2t + c_2 \sin 2t$ . Set  $Y_1 = A + Bt + Ct^2$ . Comparing the coefficients of the respective terms, we find that A = -1/8, B = 0, C = 1/4. Now set  $Y_2 = De^t$ , and obtain D = 3/5. Hence the general solution is  $y(t) = c_1 \cos 2t + c_2 \sin 2t - 1/8 + t^2/4 + 3e^t/5$ . Invoking the initial conditions, we require that  $19/40 + c_1 = 0$  and  $3/5 + 2c_2 = 2$ . Hence  $c_1 = -19/40$  and  $c_2 = 7/10$ .

13. The characteristic equation for the homogeneous problem is  $r^2 - 2r + 1 = 0$ , with a double root r = 1. Hence  $y_c(t) = c_1e^t + c_2t e^t$ . Consider  $g_1(t) = t e^t$ . Note that  $g_1$  is a solution of the homogeneous problem. Set  $Y_1 = At^2e^t + Bt^3e^t$  (the first term is not sufficient for a match). Upon substitution, we obtain  $Y_1 = t^3e^t/6$ . By inspection,  $Y_2 = 4$ . Hence the general solution is  $y(t) = c_1e^t + c_2t e^t + t^3e^t/6 + 4$ . Invoking the initial conditions, we require that  $c_1 + 4 = 1$  and  $c_1 + c_2 = 1$ . Hence  $c_1 = -3$  and  $c_2 = 4$ .

14. The characteristic equation for the homogeneous problem is  $r^2 + 4 = 0$ , with roots  $r = \pm 2i$ . Hence  $y_c(t) = c_1 \cos 2t + c_2 \sin 2t$ . Since the function  $\sin 2t$  is a solution of the homogeneous problem, set  $Y = At \cos 2t + Bt \sin 2t$ . Upon substitution, we obtain  $Y = -3t \cos 2t/4$ . Hence the general solution is  $y(t) = c_1 \cos 2t + c_2 \sin 2t - 3t \cos 2t/4$ . Invoking the initial conditions, we require that  $c_1 = 2$  and  $2c_2 - (3/4) = -1$ . Hence  $c_1 = 2$  and  $c_2 = -1/8$ .

15. The characteristic equation for the homogeneous problem is  $r^2 + 2r + 5 = 0$ , with complex roots  $r = -1 \pm 2i$ . Hence  $y_c(t) = c_1 e^{-t} \cos 2t + c_2 e^{-t} \sin 2t$ . Based on the form of g(t), set  $Y = At e^{-t} \cos 2t + Bt e^{-t} \sin 2t$ . After comparing coefficients, we obtain  $Y = t e^{-t} \sin 2t$ . Hence the general solution is  $y(t) = c_1 e^{-t} \cos 2t + c_2 e^{-t} \sin 2t + t e^{-t} \sin 2t$ . Invoking the initial conditions, we require that  $c_1 = 1$  and  $-c_1 + 2c_2 = 0$ . Hence  $c_1 = 1$  and  $c_2 = 1/2$ .

17.(a) The characteristic equation for the homogeneous problem is  $r^2 - 5r + 6 = 0$ , with roots r = 2, 3. Hence  $y_c(t) = c_1 e^{2t} + c_2 e^{3t}$ . Consider  $g_1(t) = e^{2t}(3t + 4) \sin t$ , and  $g_2(t) = e^t \cos 2t$ . Based on the form of these functions on the right hand side of the ODE, set  $Y_2(t) = e^t(A_1 \cos 2t + A_2 \sin 2t)$  and  $Y_1(t) = (B_1 + B_2 t)e^{2t} \sin t + (C_1 + C_2 t)e^{2t} \cos t$ .

(b) Substitution into the equation and comparing the coefficients results in

$$Y(t) = -\frac{1}{20}(e^t \cos 2t + 3e^t \sin 2t) + \frac{3}{2}te^{2t}(\cos t - \sin t) + e^{2t}(\frac{1}{2}\cos t - 5\sin t).$$

19.(a) The homogeneous solution is  $y_c(t) = c_1 \cos 2t + c_2 \sin 2t$ . Since  $\cos 2t$  and  $\sin 2t$  are both solutions of the homogeneous equation, set

$$Y(t) = t(A_0 + A_1t + A_2t^2)\cos 2t + t(B_0 + B_1t + B_2t^2)\sin 2t$$

(b) Substitution into the equation and comparing the coefficients results in

$$Y(t) = \left(\frac{13}{32}t - \frac{1}{12}t^3\right)\cos 2t + \frac{1}{16}(28t + 13t^2)\sin 2t.$$

20.(a) The homogeneous solution is  $y_c(t) = c_1 e^{-t} + c_2 t e^{-2t}$ . None of the functions on the right hand side are solutions of the homogeneous equation. In order to include all possible combinations of the derivatives, consider

$$Y(t) = e^{t}(A_{0} + A_{1}t + A_{2}t^{2})\cos 2t + e^{t}(B_{0} + B_{1}t + B_{2}t^{2})\sin 2t + e^{-t}(C_{1}\cos t + C_{2}\sin t) + De^{t}.$$

(b) Substitution into the differential equation and comparing the coefficients results in

$$Y(t) = e^{t}(A_{0} + A_{1}t + A_{2}t^{2})\cos 2t + e^{t}(B_{0} + B_{1}t + B_{2}t^{2})\sin 2t + e^{-t}(-\frac{3}{2}\cos t + \frac{3}{2}\sin t) + 2e^{t}/3,$$

in which  $A_0 = -4105/35152$ ,  $A_1 = 73/676$ ,  $A_2 = -5/52$ ,  $B_0 = -1233/35152$ ,  $B_1 = 10/169$ ,  $B_2 = 1/52$ .

21.(a) The homogeneous solution is  $y_c(t) = c_1 e^{-t} \cos 2t + c_2 e^{-t} \sin 2t$ . None of the terms on the right hand side are solutions of the homogeneous equation. In order to include the appropriate combinations of derivatives, consider

$$Y(t) = e^{-t}(A_1t + A_2t^2)\cos 2t + e^{-t}(B_1t + B_2t^2)\sin 2t + e^{-2t}(C_0 + C_1t)\cos 2t + e^{-2t}(D_0 + D_1t)\sin 2t.$$

(b) Substitution into the differential equation and comparing the coefficients results in

$$Y(t) = \frac{3}{16}te^{-t}\cos 2t + \frac{3}{8}t^2e^{-t}\sin 2t - \frac{1}{25}e^{-2t}(7+10t)\cos 2t + \frac{1}{25}e^{-2t}(1+5t)\sin 2t.$$

23. The homogeneous solution is  $y_c(t) = c_1 \cos \lambda t + c_2 \sin \lambda t$ . Since the differential operator does not contain a first derivative (and  $\lambda \neq m\pi$ ), we can set

$$Y(t) = \sum_{m=1}^{N} C_m \sin m\pi t \,.$$

$$-\sum_{m=1}^{N} m^2 \pi^2 C_m \sin m\pi t + \lambda^2 \sum_{m=1}^{N} C_m \sin m\pi t = \sum_{m=1}^{N} a_m \sin m\pi t.$$

Equating coefficients of the individual terms, we obtain

$$C_m = \frac{a_m}{\lambda^2 - m^2 \pi^2}, \ m = 1, 2 \dots N.$$

25. Since a, b, c > 0, the roots of the characteristic equation have negative real parts. That is,  $r = \alpha \pm \beta i$ , where  $\alpha < 0$ . Hence the homogeneous solution is

$$y_c(t) = c_1 e^{\alpha t} \cos \beta t + c_2 e^{\alpha t} \sin \beta t \,.$$

If g(t) = d, then the general solution is

$$y(t) = d/c + c_1 e^{\alpha t} \cos \beta t + c_2 e^{\alpha t} \sin \beta t.$$

Since  $\alpha < 0$ ,  $y(t) \rightarrow d/c$  as  $t \rightarrow \infty$ . If c = 0, then the characteristic roots are r = 0 and r = -b/a. The ODE becomes ay'' + by' = d. Integrating both sides, we find that  $ay' + by = dt + c_1$ . The general solution can be expressed as

$$y(t) = dt/b + c_1 + c_2 e^{-bt/a}$$

In this case, the solution grows without bound. If b = 0, also, then the differential equation can be written as y'' = d/a, which has general solution  $y(t) = dt^2/2a + c_1 + c_2$ . Hence the assertion is true only if the coefficients are positive.

27.(a) Since D is a linear operator, 
$$D^2y + bDy + cy = D^2y - (r_1 + r_2)Dy + r_1r_2y = D^2y - r_2Dy - r_1Dy + r_1r_2y = D(Dy - r_2y) - r_1(Dy - r_2y) = (D - r_1)(D - r_2)y.$$

(b) Let  $u = (D - r_2)y$ . Then the ODE (i) can be written as  $(D - r_1)u = g(t)$ , that is,  $u' - r_1u = g(t)$ . The latter is a linear first order equation in u. Its general solution is

$$u(t) = e^{r_1 t} \int_{t_0}^t e^{-r_1 \tau} g(\tau) d\tau + c_1 e^{r_1 t} d\tau$$

From above, we have  $y' - r_2 y = u(t)$ . This equation is also a first order ODE. Hence the general solution of the original second order equation is

$$y(t) = e^{r_2 t} \int_{t_0}^t e^{-r_2 \tau} u(\tau) d\tau + c_2 e^{r_2 t}.$$

Note that the solution y(t) contains two arbitrary constants.

29. We have  $(D^2 + 2D + 1)y = (D + 1)(D + 1)y$ . Let u = (D + 1)y, and consider the ODE  $u' + u = 2e^{-t}$ . The general solution is  $u(t) = 2t e^{-t} + c e^{-t}$ . We therefore have the first order equation  $u' + u = 2t e^{-t} + c_1 e^{-t}$ . The general solution of the latter differential equation is

$$y(t) = e^{-t} \int_{t_0}^t \left[2\tau + c_1\right] d\tau + c_2 e^{-t} = e^{-t} (t^2 + c_1 t + c_2).$$

30. We have  $(D^2 + 2D)y = D(D + 2)y$ . Let u = (D + 2)y, and consider the equation  $u' = 3 + 4 \sin 2t$ . Direct integration results in  $u(t) = 3t - 2 \cos 2t + c$ . The problem is reduced to solving the ODE  $y' + 2y = 3t - 2 \cos 2t + c$ . The general solution of this first order differential equation is

$$y(t) = e^{-2t} \int_{t_0}^t e^{2\tau} \left[ 3\tau - 2\cos 2\tau + c \right] d\tau + c_2 e^{-2t} =$$
$$= \frac{3}{2}t - \frac{1}{2}(\cos 2t + \sin 2t) + c_1 + c_2 e^{-2t}.$$

1. The solution of the homogeneous equation is  $y_c(t) = c_1 e^{2t} + c_2 e^{3t}$ . The functions  $y_1(t) = e^{2t}$  and  $y_2(t) = e^{3t}$  form a fundamental set of solutions. The Wronskian of these functions is  $W(y_1, y_2) = e^{5t}$ . Using the method of variation of parameters, the particular solution is given by  $Y(t) = u_1(t) y_1(t) + u_2(t) y_2(t)$ , in which

$$u_1(t) = -\int \frac{e^{3t}(2e^t)}{W(t)}dt = 2e^{-t}$$
 and  $u_2(t) = \int \frac{e^{2t}(2e^t)}{W(t)}dt = -e^{-2t}.$ 

Hence the particular solution is  $Y(t) = 2e^t - e^t = e^t$ .

3. The functions  $y_1(t) = e^{t/2}$  and  $y_2(t) = te^{t/2}$  form a fundamental set of solutions. The Wronskian of these functions is  $W(y_1, y_2) = e^t$ . First write the equation in standard form, so that  $g(t) = 4e^{t/2}$ . Using the method of variation of parameters, the particular solution is given by  $Y(t) = u_1(t) y_1(t) + u_2(t) y_2(t)$ , in which

$$u_1(t) = -\int \frac{te^{t/2}(4e^{t/2})}{W(t)}dt = -2t^2$$
 and  $u_2(t) = \int \frac{e^{t/2}(4e^{t/2})}{W(t)}dt = 4t.$ 

Hence the particular solution is  $Y(t) = -2t^2e^{t/2} + 4t^2e^{t/2} = 2t^2e^{t/2}$ .

5. The solution of the homogeneous equation is  $y_c(t) = c_1 \cos 3t + c_2 \sin 3t$ . The two functions  $y_1(t) = \cos 3t$  and  $y_2(t) = \sin 3t$  form a fundamental set of solutions, with  $W(y_1, y_2) = 3$ . The particular solution is given by  $Y(t) = u_1(t) y_1(t) + u_2(t) y_2(t)$ , in which

$$u_1(t) = -\int \frac{\sin 3t(9 \sec^2 3t)}{W(t)} dt = -\csc 3t$$
$$u_2(t) = \int \frac{\cos 3t(9 \sec^2 3t)}{W(t)} dt = \ln(\sec 3t + \tan 3t)$$

since  $0 < t < \pi/6$ . Hence  $Y(t) = -1 + (\sin 3t) \ln(\sec 3t + \tan 3t)$ . The general solution is given by

$$y(t) = c_1 \cos 3t + c_2 \sin 3t + (\sin 3t) \ln(\sec 3t + \tan 3t) - 1.$$

6. The functions  $y_1(t) = e^{-2t}$  and  $y_2(t) = te^{-2t}$  form a fundamental set of solutions. The Wronskian of these functions is  $W(y_1, y_2) = e^{-4t}$ . The particular solution is given by  $Y(t) = u_1(t) y_1(t) + u_2(t) y_2(t)$ , in which

$$u_1(t) = -\int \frac{te^{-2t}(t^{-2}e^{-2t})}{W(t)}dt = -\ln t \quad \text{and} \quad u_2(t) = \int \frac{e^{-2t}(t^{-2}e^{-2t})}{W(t)}dt = -1/t.$$

Hence the particular solution is  $Y(t) = -e^{-2t} \ln t - e^{-2t}$ . Since the second term is a solution of the homogeneous equation, the general solution is given by

$$y(t) = c_1 e^{-2t} + c_2 t e^{-2t} - e^{-2t} \ln t.$$

7. The functions  $y_1(t) = \cos(t/2)$  and  $y_2(t) = \sin(t/2)$  form a fundamental set of solutions. The Wronskian of these functions is  $W(y_1, y_2) = 1/2$ . First write the ODE in standard form, so that  $g(t) = \sec(t/2)/2$ . The particular solution is given by  $Y(t) = u_1(t) y_1(t) + u_2(t) y_2(t)$ , in which

$$u_1(t) = -\int \frac{\cos(t/2) [\sec(t/2)]}{2W(t)} dt = 2 \ln(\cos(t/2))$$
$$u_2(t) = \int \frac{\sin(t/2) [\sec(t/2)]}{2W(t)} dt = t.$$

The particular solution is  $Y(t) = 2\cos(t/2)\ln(\cos(t/2)) + t\sin(t/2)$ . The general solution is given by

$$y(t) = c_1 \cos(t/2) + c_2 \sin(t/2) + 2 \cos(t/2) \ln(\cos(t/2)) + t \sin(t/2).$$

8. The solution of the homogeneous equation is  $y_c(t) = c_1 e^t + c_2 t e^t$ . The functions  $y_1(t) = e^t$  and  $y_2(t) = t e^t$  form a fundamental set of solutions, with  $W(y_1, y_2) = e^{2t}$ . The particular solution is given by  $Y(t) = u_1(t) y_1(t) + u_2(t) y_2(t)$ , in which

$$u_1(t) = -\int \frac{te^t(e^t)}{W(t)(1+t^2)} dt = -\frac{1}{2}\ln(1+t^2)$$
$$u_2(t) = \int \frac{e^t(e^t)}{W(t)(1+t^2)} dt = \arctan t.$$

The particular solution is  $Y(t) = -(1/2)e^t \ln(1+t^2) + te^t \arctan(t)$ . Hence the general solution is given by

$$y(t) = c_1 e^t + c_2 t e^t - \frac{1}{2} e^t \ln(1+t^2) + t e^t \arctan(t).$$

10. Note first that p(t) = 0,  $q(t) = -2/t^2$  and  $g(t) = (3t^2 - 1)/t^2$ . The functions  $y_1(t)$  and  $y_2(t)$  are solutions of the homogeneous equation, verified by substitution. The Wronskian of these two functions is  $W(y_1, y_2) = -3$ . Using the method of variation of parameters, the particular solution is  $Y(t) = u_1(t) y_1(t) + u_2(t) y_2(t)$ , in which

$$u_1(t) = -\int \frac{t^{-1}(3t^2 - 1)}{t^2 W(t)} dt = t^{-2}/6 + \ln t$$

$$u_2(t) = \int \frac{t^2(3t^2 - 1)}{t^2 W(t)} dt = -t^3/3 + t/3.$$

Therefore  $Y(t) = 1/6 + t^2 \ln t - t^2/3 + 1/3$ .

12. Observe that  $g(t) = t e^{2t}$ . The functions  $y_1(t)$  and  $y_2(t)$  are a fundamental set of solutions. The Wronskian of these two functions is  $W(y_1, y_2) = t e^t$ . Using the method of variation of parameters, the particular solution is  $Y(t) = u_1(t) y_1(t) + u_2(t) y_2(t)$ , in which

$$u_1(t) = -\int \frac{e^t(t e^{2t})}{W(t)} dt = -e^{2t}/2 \quad \text{and} \quad u_2(t) = \int \frac{(1+t)(t e^{2t})}{W(t)} dt = t e^t.$$
  
Therefore  $Y(t) = -(1+t)e^{2t}/2 + t e^{2t} = -e^{2t}/2 + t e^{2t}/2.$ 

13. Note that  $g(x) = \ln x$ . The functions  $y_1(x) = x^2$  and  $y_2(x) = x^2 \ln x$  are solutions of the homogeneous equation, as verified by substitution. The Wronskian of the solutions is  $W(y_1,y_2) = x^3$ . Using the method of variation of parameters, the particular solution is  $Y(x) = u_1(x) y_1(x) + u_2(x) y_2(x)$ , in which

$$u_1(x) = -\int \frac{x^2 \ln x(\ln x)}{W(x)} dx = -(\ln x)^3/3$$
$$u_2(x) = \int \frac{x^2(\ln x)}{W(x)} dx = (\ln x)^2/2.$$

Therefore  $Y(x) = -x^2(\ln x)^3/3 + x^2(\ln x)^3/2 = x^2(\ln x)^3/6$ .

15. First write the equation in standard form. The forcing function becomes  $g(x)/x^2$ . The functions  $y_1(x) = x^{-1/2} \sin x$  and  $y_2(x) = x^{-1/2} \cos x$  are a fundamental set of solutions. The Wronskian of the solutions is  $W(y_1,y_2) = -1/x$ . Using the method of variation of parameters, the particular solution is  $Y(x) = u_1(x) y_1(x) + u_2(x) y_2(x)$ , in which

$$u_1(x) = \int_{x_0}^x \frac{\cos \tau \left(g(\tau)\right)}{\tau \sqrt{\tau}} d\tau \quad \text{and} \quad u_2(x) = -\int_{x_0}^x \frac{\sin \tau \left(g(\tau)\right)}{\tau \sqrt{\tau}} d\tau.$$

Therefore

$$Y(x) = \frac{\sin x}{\sqrt{x}} \int_{x_0}^x \frac{\cos \tau \left(g(\tau)\right)}{\tau \sqrt{\tau}} dt - \frac{\cos x}{\sqrt{x}} \int_{x_0}^x \frac{\sin \tau \left(g(\tau)\right)}{\tau \sqrt{\tau}} d\tau =$$
$$= \frac{1}{\sqrt{x}} \int_{x_0}^x \frac{\sin(x-\tau) g(\tau)}{\tau \sqrt{\tau}} d\tau.$$

16. Eq.(28) is

$$Y(t) = -y_1(t) \int_{t_0}^t \frac{y_2(s)g(s)}{W(y_1, y_2)(s)} \, ds + y_2(t) \int_{t_0}^t \frac{y_1(s)g(s)}{W(y_1, y_2)(s)} \, ds$$

where  $t_0$  is now considered the initial point. Bringing the terms  $y_1(t)$  and  $y_2(t)$  inside the integrals and using the fact that  $W(y_1, y_2)(s) = y_1(s)y'_2(s) - y'_1(s)y_2(s)$ , the desired result holds. To show that Y(t) satisfies L[y] = g(t) we must take the derivative using Leibniz's rule, which says that if  $y(t) = \int_{t_0}^t G(t, s) ds$ , then

 $Y'(t) = G(t,t) + \int_{t_0}^t G_t(t,s) \, ds$ . Letting G(t,s) be the above integrand, we have that G(t,t) = 0 and

$$\frac{\partial G}{\partial t} = \frac{y_1(s)y_2'(t) - y_1'(t)y_2(s)}{W(y_1, y_2)(s)}g(s).$$

Likewise,

$$Y'' = \frac{\partial G(t,t)}{\partial t} + \int_{t_0}^t \frac{\partial^2 G}{\partial t^2}(t,s) \, ds = g(t) + \int_{t_0}^t \frac{y_1(s)y_2''(t) - y_1''(t)y_2(s)}{W(y_1,y_2)(s)}g(s) \, ds.$$

Since  $y_1$  and  $y_2$  are solutions of L[y] = 0, we have L[Y] = g(t) since all the terms involving the integral will add to zero. Clearly  $y(t_0) = 0$  and  $y'(t_0) = 0$ .

17. Let  $y_1(t)$  and  $y_2(t)$  be a fundamental set of solutions, and  $W(t) = W(y_1, y_2)$  be the corresponding Wronskian. Any solution, u(t), of the homogeneous equation is a linear combination  $u(t) = \alpha_1 y_1(t) + \alpha_2 y_2(t)$ . Invoking the initial conditions, we require that

$$egin{aligned} y_0 &= lpha_1 \, y_1(t_0) + lpha_2 \, y_2(t_0) \ y_0' &= lpha_1 \, y_1'(t_0) + lpha_2 \, y_2'(t_0) \end{aligned}$$

Note that this system of equations has a unique solution, since  $W(t_0) \neq 0$ . Now consider the nonhomogeneous problem, L[v] = g(t), with homogeneous initial conditions. Using the method of variation of parameters, the particular solution is given by

$$Y(t) = -y_1(t) \int_{t_0}^t \frac{y_2(s) g(s)}{W(s)} ds + y_2(t) \int_{t_0}^t \frac{y_1(s) g(s)}{W(s)} ds \, .$$

The general solution of the IVP (iii) is

$$v(t) = \beta_1 y_1(t) + \beta_2 y_2(t) + Y(t) = \beta_1 y_1(t) + \beta_2 y_2(t) + y_1(t)u_1(t) + y_2(t)u_2(t)$$

in which  $u_1$  and  $u_2$  are defined above. Invoking the initial conditions, we require that

$$0 = \beta_1 y_1(t_0) + \beta_2 y_2(t_0) + Y(t_0)$$
  
$$0 = \beta_1 y_1'(t_0) + \beta_2 y_2'(t_0) + Y'(t_0)$$

Based on the definition of  $u_1$  and  $u_2$ ,  $Y(t_0) = 0$ . Furthermore, since  $y_1u'_1 + y_2u'_2 = 0$ , it follows that  $Y'(t_0) = 0$ . Hence the only solution of the above system of equations is the trivial solution. Therefore v(t) = Y(t). Now consider the function y = u + v. Then L[y] = L[u + v] = L[u] + L[v] = g(t). That is, y(t) is a solution of the nonhomogeneous problem. Further,  $y(t_0) = u(t_0) + v(t_0) = y_0$ , and similarly,  $y'(t_0) = y'_0$ . By the uniqueness theorems, y(t) is the unique solution of the initial value problem.

18.(a) A fundamental set of solutions is  $y_1(t) = \cos t$  and  $y_2(t) = \sin t$ . The Wronskian  $W(t) = y_1y_2' - y_1'y_2 = 1$ . By the result in Problem 17,

$$Y(t) = \int_{t_0}^t \frac{\cos(s)\,\sin(t) - \cos(t)\,\sin(s)}{W(s)} g(s)ds$$
  
=  $\int_{t_0}^t \left[\cos(s)\,\sin(t) - \cos(t)\,\sin(s)\right] g(s)ds$ .

Finally, we have  $\cos(s) \sin(t) - \cos(t) \sin(s) = \sin(t - s)$ .

(b) Using Problem 16 and part (a), the solution is

$$y(t) = y_0 \cos t + y'_0 \sin t + \int_0^t \sin(t-s)g(s)ds$$

19. A fundamental set of solutions is  $y_1(t) = e^{at}$  and  $y_2(t) = e^{bt}$ . The Wronskian  $W(t) = y_1y'_2 - y'_1y_2 = (b-a)e^{(a+b)t}$ . By the result in Problem 17,

$$Y(t) = \int_{t_0}^t \frac{e^{as}e^{bt} - e^{at}e^{bs}}{W(s)}g(s)ds = \frac{1}{b-a}\int_{t_0}^t \frac{e^{as}e^{bt} - e^{at}e^{bs}}{e^{(a+b)s}}g(s)ds = \frac{1}{b-a}\int_{t_0}^t \frac{e^{as}e^{bt} - e^{at}e^{bs}}{e^{(a+b)s}}g(s)ds$$

Hence the particular solution is

$$Y(t) = \frac{1}{b-a} \int_{t_0}^t \left[ e^{b(t-s)} - e^{a(t-s)} \right] g(s) ds$$

21. A fundamental set of solutions is  $y_1(t) = e^{at}$  and  $y_2(t) = te^{at}$ . The Wronskian  $W(t) = y_1y'_2 - y'_1y_2 = e^{2at}$ . By the result in Problem 17,

$$Y(t) = \int_{t_0}^t \frac{te^{as+at} - s e^{at+as}}{W(s)} g(s) ds = \int_{t_0}^t \frac{(t-s)e^{as+at}}{e^{2as}} g(s) ds$$

Hence the particular solution is

$$Y(t) = \int_{t_0}^t (t-s)e^{a(t-s)}g(s)ds \,.$$

22. The form of the kernel depends on the characteristic roots. If the roots are real and distinct,

$$K(t-s) = \frac{e^{b(t-s)} - e^{a(t-s)}}{b-a}.$$

If the roots are real and identical,

$$K(t-s) = (t-s)e^{a(t-s)}$$

If the roots are complex conjugates,

$$K(t-s) = \frac{e^{\lambda(t-s)} \sin \mu(t-s)}{\mu}.$$

23. Let  $y(t) = v(t)y_1(t)$ , in which  $y_1(t)$  is a solution of the homogeneous equation. Substitution into the given ODE results in

$$v''y_1 + 2v'y_1' + vy_1'' + p(t)[v'y_1 + vy_1'] + q(t)vy_1 = g(t)$$

By assumption,  $y_1'' + p(t)y_1 + q(t)y_1 = 0$ , hence v(t) must be a solution of the ODE

$$v''y_1 + [2y_1' + p(t)y_1]v' = g(t)$$

Setting w = v', we also have  $w'y_1 + [2y'_1 + p(t)y_1]w = g(t)$ .

25. First write the equation as  $y'' + 7t^{-1}y + 5t^{-2}y = t^{-1}$ . As shown in Problem 23, the function  $y(t) = t^{-1}v(t)$  is a solution of the given ODE as long as v is a solution of

$$t^{-1} v'' + \left[-2t^{-2} + 7t^{-2}\right] v' = t^{-1},$$

that is,  $v'' + 5t^{-1}v' = 1$ . This ODE is linear and first order in v'. The integrating factor is  $\mu = t^5$ . The solution is  $v' = t/6 + ct^{-5}$ . Direct integration now results in  $v(t) = t^2/12 + c_1t^{-4} + c_2$ . Hence  $y(t) = t/12 + c_1t^{-5} + c_2t^{-1}$ .

26. Write the equation as  $y'' - t^{-1}(1+t)y + t^{-1}y = t e^{2t}$ . As shown in Problem 23, the function y(t) = (1+t)v(t) is a solution of the given ODE as long as v is a solution of

$$(1+t)v'' + \left[2 - t^{-1}(1+t)^2\right]v' = t e^{2t},$$

that is,

$$v'' - \frac{1+t^2}{t(t+1)}v' = \frac{t}{t+1}e^{2t}.$$

This equation is first order linear in v', with integrating factor  $\mu = t^{-1}(1+t)^2 e^{-t}$ . The solution is  $v' = (t^2 e^{2t} + c_1 t e^t)/(1+t)^2$ . Integrating, we obtain  $v(t) = e^{2t}/2 - e^{2t}/(t+1) + c_1 e^t/(t+1) + c_2$ . Hence the solution of the original ODE is  $y(t) = (t-1)e^{2t}/2 + c_1 e^t + c_2(t+1)$ .

1.  $R \cos \delta = 3$  and  $R \sin \delta = 4$ , so  $R = \sqrt{25} = 5$  and  $\delta = \arctan(4/3)$ . We obtain that  $u = 5 \cos(2t - \arctan(4/3))$ .

2.  $R \cos \delta = -2$  and  $R \sin \delta = -3$ , so  $R = \sqrt{13}$  and  $\delta = \pi + \arctan(3/2)$ . We obtain that  $u = \sqrt{13} \cos(\pi t - \pi - \arctan(3/2))$ .

4. The spring constant is k = 3/(1/4) = 12 lb/ft. Mass m = 3/32 lb-s<sup>2</sup>/ft. Since there is no damping, the equation of motion is 3u''/32 + 12u = 0, that is, u'' + 128u = 0. The initial conditions are u(0) = -1/12 ft, u'(0) = 2 ft/s. The general solution is  $u(t) = A \cos 8\sqrt{2}t + B \sin 8\sqrt{2}t$ . Invoking the initial conditions, we have

$$u(t) = -\frac{1}{12} \cos 8\sqrt{2}t + \frac{1}{4\sqrt{2}} \sin 8\sqrt{2}t.$$

 $R = \sqrt{11/288}$  ft,  $\delta = \pi - \arctan(3/\sqrt{2})$  rad,  $\omega_0 = 8\sqrt{2}$  rad/s,  $T = \pi/(4\sqrt{2})$  s.

6. The spring constant is k = 3/(.1) = 30 N/m. The damping coefficient is given as  $\gamma = 3/5$  N-s/m. Hence the equation of motion is 2u'' + 3u'/5 + 30u = 0, that is, u'' + 0.3u' + 15u = 0. The initial conditions are u(0) = 0.05 m and u'(0) = 0.01 m/s. The general solution is  $u(t) = A \cos \mu t + B \sin \mu t$ , in which  $\mu = 3.87008$  rad/s. Invoking the initial conditions, we have  $u(t) = e^{-0.15t}(0.05 \cos \mu t + 0.00452 \sin \mu t)$ . Also,  $\mu/\omega_0 = 3.87008/\sqrt{15} \approx 0.99925$ .

3.7

8. The frequency of the undamped motion is  $\omega_0 = 1$ . The quasi frequency of the damped motion is  $\mu = \sqrt{4 - \gamma^2}/2$ . Setting  $\mu = 2\omega_0/3$ , we obtain  $\gamma = 2\sqrt{5}/3$ .

9. The spring constant is k = mg/L. The equation of motion for an undamped system is mu'' + mgu/L = 0. Hence the natural frequency of the system is  $\omega_0 = \sqrt{g/L}$ . The period is  $T = 2\pi/\omega_0$ .

10. The general solution of the system is  $u(t) = A \cos \gamma(t - t_0) + B \sin \gamma(t - t_0)$ . Invoking the initial conditions, we have  $u(t) = u_0 \cos \gamma(t - t_0) + (u'_0/\gamma) \sin \gamma(t - t_0)$ . Clearly, the functions  $v = u_0 \cos \gamma(t - t_0)$  and  $w = (u'_0/\gamma) \sin \gamma(t - t_0)$  satisfy the given criteria.

11. Note that  $r \sin(\omega_0 t - \theta) = r \sin \omega_0 t \cos \theta - r \cos \omega_0 t \sin \theta$ . Comparing the given expressions, we have  $A = -r \sin \theta$  and  $B = r \cos \theta$ . That is,  $r = R = \sqrt{A^2 + B^2}$ , and  $\tan \theta = -A/B = -1/\tan \delta$ . The latter relation is also  $\tan \theta + \cot \delta = 1$ .

12. The system is critically damped, when  $R = 2\sqrt{L/C}$ . Here R = 1000 ohms.

15.(a) Let  $u = Re^{-\gamma t/2m} \cos(\mu t - \delta)$ . Then attains a maximum when  $\mu t_k - \delta = 2k\pi$ . Hence  $T_d = t_{k+1} - t_k = 2\pi/\mu$ .

(b)  $u(t_k)/u(t_{k+1}) = e^{-\gamma t_k/2m}/e^{-\gamma t_{k+1}/2m} = e^{(\gamma t_{k+1} - \gamma t_k)/2m}$ . Hence  $u(t_k)/u(t_{k+1}) = e^{\gamma(2\pi/\mu)/2m} = e^{\gamma T_d/2m}$ .

(c) 
$$\Delta = \ln \left[ u(t_k) / u(t_{k+1}) \right] = \gamma (2\pi/\mu) / 2m = \pi \gamma / \mu m$$
.

16. The spring constant is k = 16/(1/4) = 64 lb/ft. Mass m = 1/2 lb-s<sup>2</sup>/ft. The damping coefficient is  $\gamma = 2$  lb-s/ft. The quasi frequency is  $\mu = 2\sqrt{31}$  rad/s. Hence  $\Delta = 2\pi/\sqrt{31} \approx 1.1285$ .

18.(a) The characteristic equation is  $mr^2 + \gamma r + k = 0$ . Since  $\gamma^2 < 4km$ , the roots are  $r_{1,2} = (-\gamma \pm i\sqrt{4mk - \gamma^2})/2m$ . The general solution is

$$u(t) = e^{-\gamma t/2m} \left[ A \cos \frac{\sqrt{4mk - \gamma^2}}{2m} t + B \sin \frac{\sqrt{4mk - \gamma^2}}{2m} t \right]$$

Invoking the initial conditions,  $A = u_0$  and  $B = (2mv_0 - \gamma u_0)/\sqrt{4mk - \gamma^2}$ .

(b) We can write  $u(t) = R e^{-\gamma t/2m} \cos(\mu t - \delta)$ , in which

$$R = \sqrt{u_0^2 + \frac{(2mv_0 - \gamma u_0)^2}{4mk - \gamma^2}} \quad \text{and} \quad \delta = \arctan\left[\frac{(2mv_0 - \gamma u_0)}{u_0\sqrt{4mk - \gamma^2}}\right].$$

(c)

$$R = \sqrt{u_0^2 + \frac{(2mv_0 - \gamma u_0)^2}{4mk - \gamma^2}} = 2\sqrt{\frac{m(ku_0^2 + \gamma u_0v_0 + mv_0^2)}{4mk - \gamma^2}} = \sqrt{\frac{a + b\gamma}{4mk - \gamma^2}}$$

It is evident that R increases (monotonically) without bound as  $\gamma \rightarrow (2\sqrt{mk})^-$ .

20.(a) The general solution is  $u(t) = A \cos \sqrt{2} t + B \sin \sqrt{2} t$ . Invoking the initial conditions, we have  $u(t) = \sqrt{2} \sin \sqrt{2} t$ .





The condition u'(0) = 2 implies that u(t) initially increases. Hence the phase point travels clockwise.

23. Based on Newton's second law, with the positive direction to the right,  $\sum F = mu''$ , where  $\sum F = -ku - \gamma u'$ . Hence the equation of motion is  $mu'' + \gamma u' + ku = 0$ . The only difference in this problem is that the equilibrium position is located at the unstretched configuration of the spring.

24.(a) The restoring force exerted by the spring is  $F_s = -(ku + \epsilon u^3)$ . The opposing viscous force is  $F_d = -\gamma u'$ . Based on Newton's second law, with the positive direction to the right,  $F_s + F_d = mu''$ . Hence the equation of motion is  $mu'' + \gamma u' + ku + \epsilon u^3 = 0$ .

(b) With the specified parameter values, the equation of motion is u'' + u = 0. The general solution of this ODE is  $u(t) = A \cos t + B \sin t$ . Invoking the initial conditions, the specific solution is  $u(t) = \sin t$ . Clearly, the amplitude is R = 1, and the period of the motion is  $T = 2\pi$ . (c) Given  $\epsilon = 0.1$ , the equation of motion is  $u'' + u + 0.1 u^3 = 0$ . A solution of the IVP can be generated numerically. We estimate A = 0.98 and T = 6.07.



(d) For  $\epsilon = 0.2$ , A = 0.96 and T = 5.90. For  $\epsilon = 0.3$ , A = 0.94 and T = 5.74.



(e) The amplitude and period both seem to decrease.

(f) For  $\epsilon = -0.1$ , A = 1.03 and T = 6.55. For  $\epsilon = -0.2$ , A = 1.06 and T = 6.90. For  $\epsilon = -0.3$ , A = 1.11 and T = 7.41. The amplitude and period both seem to increase.



3.8

1. We have  $\sin(\alpha \pm \beta) = \sin \alpha \cos \beta \pm \cos \alpha \sin \beta$ . Subtracting the two identities, we obtain  $\sin(\alpha + \beta) - \sin(\alpha - \beta) = 2 \cos \alpha \sin \beta$ . Setting  $\alpha + \beta = 7t$  and  $\alpha - \beta = 6t$ , we get that  $\alpha = 6.5t$  and  $\beta = 0.5t$ . This implies that  $\sin 7t - \sin 6t = 2 \sin (t/2) \cos (13t/2)$ .

2. Consider the trigonometric identities  $\cos(\alpha \pm \beta) = \cos \alpha \cos \beta \mp \sin \alpha \sin \beta$ . Adding the two identities, we get  $\cos(\alpha - \beta) + \cos(\alpha + \beta) = 2 \cos \alpha \cos \beta$ . Comparing the expressions, set  $\alpha + \beta = 2\pi t$  and  $\alpha - \beta = \pi t$ . This means  $\alpha = 3\pi t/2$  and  $\beta = \pi t/2$ . Upon substitution, we have  $\cos(\pi t) + \cos(2\pi t) = 2 \cos(3\pi t/2) \cos(\pi t/2)$ .

3. Adding the two identities  $\sin(\alpha \pm \beta) = \sin \alpha \cos \beta \pm \cos \alpha \sin \beta$ , it follows that  $\sin(\alpha - \beta) + \sin(\alpha + \beta) = 2 \sin \alpha \cos \beta$ . Setting  $\alpha + \beta = 4t$  and  $\alpha - \beta = 3t$ , we have  $\alpha = 7t/2$  and  $\beta = t/2$ . Hence  $\sin 3t + \sin 4t = 2 \sin(7t/2) \cos(t/2)$ .

4. Using MKS units, the spring constant is k = 5(9.8)/0.1 = 490 N/m, and the damping coefficient is  $\gamma = 2/0.04 = 50$  N-s/m. The equation of motion is

$$5u'' + 50u' + 490u = 10\sin(t/2).$$

The initial conditions are u(0) = 0 m and u'(0) = 0.03 m/s.

5.(a) The homogeneous solution is  $u_c(t) = Ae^{-5t} \cos \sqrt{73} t + Be^{-5t} \sin \sqrt{73} t$ . Based on the method of undetermined coefficients, the particular solution is

$$U(t) = \frac{1}{153281} \left[ -160 \cos(t/2) + 3128 \sin(t/2) \right].$$

Hence the general solution of the ODE is  $u(t) = u_c(t) + U(t)$ . Invoking the initial conditions, we find that

$$A = 160/153281$$
 and  $B = 383443\sqrt{73}/1118951300$ .

Hence the response is

$$u(t) = \frac{1}{153281} \left[ 160 \, e^{-5t} \cos \sqrt{73} \, t + \frac{383443\sqrt{73}}{7300} e^{-5t} \sin \sqrt{73} \, t \right] + U(t).$$

(b)  $u_c(t)$  is the transient part and U(t) is the steady state part of the response.





(d) The amplitude of the forced response is given by  $R = 2/\Delta$ , in which

$$\Delta = \sqrt{25(98 - \omega^2)^2 + 2500\,\omega^2}$$

The maximum amplitude is attained when  $\Delta$  is a minimum. Hence the amplitude is maximum at  $\omega = 4\sqrt{3}$  rad/s.

8. The equation of motion is  $2u'' + u' + 3u = 3\cos 3t - 2\sin 3t$ . Since the system is damped, the steady state response is equal to the particular solution. Using the method of undetermined coefficients, we obtain  $u_{ss}(t) = (\sin 3t - \cos 3t)/6$ . Further, we find that  $R = \sqrt{2}/6$  and  $\delta = \arctan(-1) = 3\pi/4$ . Hence we can write  $u_{ss}(t) = (\sqrt{2}/6)\cos(3t - 3\pi/4)$ .

9.(a) Plug in  $u(t) = R\cos(\omega t - \delta)$  into the equation  $mu'' + \gamma u' + ku = F_0 \cos \omega t$ , then use trigonometric identities and compare the coefficients of  $\cos \omega t$  and  $\sin \omega t$ . The result follows.

(b) First note that since  $R = F_0/\Delta$ ,  $Rk/F_0 = k/\Delta$  and that since  $\Gamma = \gamma^2/(mk)$ ,  $(\gamma^2 \omega^2)/m^2 = \Gamma \omega_0^2 \omega^2$ . Then using Eq.12,

$$\begin{aligned} \frac{Rk}{F_0} &= \frac{k}{\Delta} = \frac{m\omega_0^2}{\sqrt{m^2(\omega_0^2 - \omega^2)^2 + \gamma^2 \omega^2}} = \frac{m\omega_0^2}{\sqrt{m^2(\omega_0^2 - \omega^2)^2 + \gamma^2 \omega^2}} \\ &= \frac{\omega_0^2}{\sqrt{(\omega_0^2 - \omega^2)^2 + \frac{\gamma^2 \omega^2}{m^2}}} = \frac{\omega_0^2}{\sqrt{(\omega_0^2 - \omega^2)^2 + \Gamma \omega_0^2 \omega^2}} \\ &= \frac{1}{\sqrt{\left(\frac{\omega_0^2 - \omega^2}{\omega_0^2}\right)^2 + \Gamma \frac{\omega_0^2 \omega^2}{\omega_0^4}}} = \frac{1}{\sqrt{\left(1 - \frac{\omega^2}{\omega_0^2}\right)^2 + \Gamma \frac{\omega^2}{\omega_0^2}}} \end{aligned}$$

(c) The amplitude of the steady-state response is given by

$$R = \frac{F_0}{\sqrt{m^2(\omega_0^2 - \omega^2)^2 + \gamma^2 \,\omega^2}}$$

Since  $F_0$  is constant, the amplitude is maximum when the denominator of R is minimum. Let  $z = \omega^2$ , and consider the function  $f(z) = m^2(\omega_0^2 - z)^2 + \gamma^2 z$ . Note

that f(z) is a quadratic, with minimum at  $z = \omega_0^2 - \gamma^2/2m^2$ . Hence the amplitude R attains a maximum at  $\omega_{max}^2 = \omega_0^2 - \gamma^2/2m^2$ . Furthermore, since  $\omega_0^2 = k/m$ ,

$$\omega_{max}^2 = \omega_0^2 \left[ 1 - \frac{\gamma^2}{2km} \right].$$

(d) Substituting  $\omega^2 = \omega_{max}^2$  into the expression for the amplitude R gives the maximum value for R:

$$R_{max} = \frac{F_0}{\sqrt{\gamma^4/4m^2 + \gamma^2 \left(\omega_0^2 - \gamma^2/2m^2\right)}} = \frac{F_0}{\sqrt{\omega_0^2 \gamma^2 - \gamma^4/4m^2}} = \frac{F_0}{\gamma \omega_0 \sqrt{1 - \gamma^2/4mk}}$$

To understand the approximation, note that

$$R_{max} = \frac{F_0}{\gamma\omega_0} \left(1 - \frac{\gamma^2}{4mk}\right)^{-1/2}$$

Recall that binomial theorem states that  $(1 + a)^p \approx 1 + pa$  when a is small. Applying this result with  $a = -\gamma^2/(4mk)$  and p = -1/2 gives that

$$R_{max} = \frac{F_0}{\gamma\omega_0} \left(1 - \frac{\gamma^2}{4mk}\right)^{-1/2} \approx \frac{F_0}{\gamma\omega_0} \left(1 + \left(-\frac{1}{2}\right)\left(-\frac{\gamma^2}{4mk}\right)\right) = \frac{F_0}{\gamma\omega_0} \left(1 + \frac{\gamma^2}{8mk}\right)$$

13.(a) The homogeneous solution is  $u_c(t) = A \cos t + B \sin t$ . Based on the method of undetermined coefficients, the particular solution is

$$U(t) = \frac{3}{1 - \omega^2} \cos \omega t \,.$$

Hence the general solution of the ODE is  $u(t) = u_c(t) + U(t)$ . Invoking the initial conditions, we find that  $A = 3/(\omega^2 - 1)$  and B = 0. Hence the response is

$$u(t) = \frac{3}{1 - \omega^2} \left[ \cos \omega t - \cos t \right].$$

(b)

$$(a) \ \omega = 0.7 \qquad (b) \ \omega = 0.8 \qquad (c) \ \omega = 0.9$$

Note that

$$u(t) = \frac{6}{1 - \omega^2} \sin\left[\frac{(1 - \omega)t}{2}\right] \sin\left[\frac{(\omega + 1)t}{2}\right].$$

14.(a) The homogeneous solution is  $u_c(t) = A \cos t + B \sin t$ . Based on the method of undetermined coefficients, the particular solution is

$$U(t) = \frac{3}{1 - \omega^2} \cos \omega t \,.$$

Hence the general solution is  $u(t) = u_c(t) + U(t)$ . Invoking the initial conditions, we find that  $A = (\omega^2 + 2)/(\omega^2 - 1)$  and B = 1. Hence the response is

$$u(t) = \frac{1}{1 - \omega^2} \left[ 3 \cos \omega t - (\omega^2 + 2) \cos t \right] + \sin t.$$

(b)



Note that

$$u(t) = \frac{6}{1 - \omega^2} \sin\left[\frac{(1 - \omega)t}{2}\right] \sin\left[\frac{(\omega + 1)t}{2}\right] + \cos t + \sin t.$$

15.



18.(a)



(b) Phase plot - u' vs u:



### **Elementary Differential Equations 11th Edition Boyce Solutions Manual**

Full Download: https://alibabadownload.com/product/elementary-differential-equations-11th-edition-boyce-solutions-manual/

84

Chapter 3. Second-Order Linear Equations