Full Download: https://alibabadownload.com/product/statistics-for-managers-using-ms-excel-global-7th-edition-levine-solutions-managers-using-ms-excel-global-7th-edition-levine-solutions-managers-using-ms-excel-global-7th-edition-levine-solutions-managers-using-ms-excel-global-7th-edition-levine-solutions-managers-using-ms-excel-global-7th-edition-levine-solutions-managers-using-ms-excel-global-7th-edition-levine-solutions-managers-using-ms-excel-global-7th-edition-levine-solutions-managers-using-ms-excel-global-7th-edition-levine-solutions-managers-using-ms-excel-global-7th-edition-levine-solutions-managers-using-ms-excel-global-7th-edition-levine-solutions-managers-using-ms-excel-global-7th-edition-levine-solutions-managers-using-ms-excel-global-7th-edition-levine-solutions-managers-using-ms-excel-global-7th-edition-levine-solutions-managers-using-ms-excel-global-7th-edition-levine-solutions-managers-using-ms-excel-global-7th-edition-levine-solutions-managers-managers-using-ms-excel-global-7th-edition-levine-solutions-managers-managers-using-ms-excel-global-7th-edition-levine-solutions-managers-managers-using-ms-excel-global-7th-edition-levine-solutions-managers-

38 Chapter 1: Defining and Collecting Data

CHAPTER 1

- 1.1 (a) The type of beverage sold yields categorical or "qualitative" responses.
 - (b) The type of beverage sold yields distinct categories in which no ordering is implied.
- 1.2 Three sizes of U.S. businesses are classified into distinct categories—small, medium, and large—in which order is implied.
- 1.3 (a) The time it takes to download a video from the Internet is a continuous numerical or "quantitative" variable because time can have any value from 0 to any reasonable unit of time.
 - (b) The download time is a ratio scaled variable because the true zero point in the measurement is zero units of time.
- 1.4 (a) The number of cellphones is a numerical variable that is discrete because the outcome is a count. It is ratio scaled because it has a true zero point.
 - (b) Monthly data usage is a numerical variable that is continuous because any value within a range of values can occur. It is ratio scaled because it has a true zero point.
 - (c) Number of text messages exchanged per month is a numerical variable that is discrete because the outcome is a count. It is ratio scaled because it has a true zero point.
 - (d) Voice usage per month is a numerical variable that is continuous because any value within a range of values can occur. It is ratio scaled because it has a true zero point.
 - (e) Whether a cellphone is used for email is a categorical variable because the answer can be only yes or no. This also makes it a nominal-scaled variable.
- 1.5 (a) numerical, continuous, ratio scale
 - (b) numerical, discrete, ratio scale
 - (c) categorical, nominal scale
 - (d) categorical, nominal scale
- 1.6 (a) Categorical, nominal scale.
 - (b) Numerical, continuous, ratio scale.
 - (c) Categorical, nominal scale.
 - (d) Numerical, discrete, ratio scale.
 - (e) Categorical, nominal scale.
- 1.7 (a) numerical, continuous, ratio scale *
 - (b) categorical, nominal scale
 - (c) categorical, nominal scale
 - (d) numerical, discrete, ratio scale

*Some researchers consider money as a discrete numerical variable because it can be "counted."

- 1.8 (a) numerical, continuous, ratio scale *
 - (b) numerical, discrete, ratio scale
 - (c) numerical, continuous, ratio scale *
 - (d) categorical, nominal

*Some researchers consider money as a discrete numerical variable because it can be "counted."

- 1.9 (a) Income may be considered discrete if we "count" our money. It may be considered continuous if we "measure" our money; we are only limited by the way a country's monetary system treats its currency.
 - (b) The first format is preferred because the responses represent data measured on a higher scale.
- 1.10 The underlying variable, ability of the students, may be continuous, but the measuring device, the test, does not have enough precision to distinguish between the two students.
- 1.11 (a) The population is "all working women from the metropolitan area." A systematic or random sample could be taken of women from the metropolitan area. The director might wish to collect both numerical and categorical data.
 - (b) Three categorical questions might be occupation, marital status, type of clothing. Numerical questions might be age, average monthly hours shopping for clothing, income.
- 1.12 The answer depends on the chosen data set.
- 1.13 The answer depends on the specific story.
- 1.14 The answer depends on the specific story.
- 1.15 The transportation engineers and planners should use primary data collected through an observational study of the driving characteristics of drivers over the course of a month.
- 1.16 The information presented there is based on surveys.
- 1.17 (a) 001 (b) 040 (c) 902
- 1.18 Sample without replacement: Read from left to right in 3-digit sequences and continue unfinished sequences from end of row to beginning of next row.
 Row 05: 338 505 855 551 438 855 077 186 579 488 767 833 170
 Rows 05-06: 897
 Row 06: 340 033 648 847 204 334 639 193 639 411 095 924
 Rows 06-07: 707
 Row 07: 054 329 776 100 871 007 255 980 646 886 823 920 461
 Row 08: 893 829 380 900 796 959 453 410 181 277 660 908 887
 Rows 08-09: 237
 Row 09: 818 721 426 714 050 785 223 801 670 353 362 449
 Rows 09-10: 406
 Note: All sequences above 902 and duplicates are discarded.

40 Chapter 1: Defining and Collecting Data

- 1.19
 (a)
 Row 29: 12 47 83 76 22 99 65 93 10 65 83 61 36 98 89 58 86 92 71

 Note: All sequences above 93 and all repeating sequences are discarded.
 - (b) Row 29: 12 47 83 76 22 99 65 93 10 65 83 61 36 98 89 58 86 Note: All sequences above 93 are discarded. Elements 65 and 83 are repeated.
- 1.20 A simple random sample would be less practical for personal interviews because of travel costs (unless interviewees are paid to attend a central interviewing location).
- 1.21 This is a probability sample because the selection is based on chance. It is not a simple random sample because A is more likely to be selected than B or C.
- 1.22 Here all members of the population are equally likely to be selected and the sample selection mechanism is based on chance. But not every sample of size 2 has the same chance of being selected. For example the sample "B and C" is impossible.
- 1.23 (a) Since a complete roster of full-time students exists, a simple random sample of 200 students could be taken. If student satisfaction with the quality of campus life randomly fluctuates across the student body, a systematic 1-in-20 sample could also be taken from the population frame. If student satisfaction with the quality of life may differ by gender and by experience/class level, a stratified sample using eight strata, female freshmen through female seniors and male freshmen through male seniors, could be selected. If student satisfaction with the quality of life is thought to fluctuate as much within clusters as between them, a cluster sample could be taken.
 - (b) A simple random sample is one of the simplest to select. The population frame is the registrar's file of 4,000 student names.
 - (c) A systematic sample is easier to select by hand from the registrar's records than a simple random sample, since an initial person at random is selected and then every 20th person thereafter would be sampled. The systematic sample would have the additional benefit that the alphabetic distribution of sampled students' names would be more comparable to the alphabetic distribution of student names in the campus population.
 - (d) If rosters by gender and class designations are readily available, a stratified sample should be taken. Since student satisfaction with the quality of life may indeed differ by gender and class level, the use of a stratified sampling design will not only ensure all strata are represented in the sample, it will also generate a more representative sample and produce estimates of the population parameter that have greater precision.
 - (e) If all 4,000 full-time students reside in one of 10 on-campus residence halls which fully integrate students by gender and by class, a cluster sample should be taken. A cluster could be defined as an entire residence hall, and the students of a single randomly selected residence hall could be sampled. Since each dormitory has 400 students, a systematic sample of 200 students can then be selected from the chosen cluster of 400 students. Alternately, a cluster could be defined as a floor of one of the 10 dormitories. Suppose there are four floors in each dormitory with 100 students on each floor. Two floors could be randomly sampled to produce the required 200 student sample. Selection of an entire dormitory may make distribution and collection of the survey easier to accomplish. In contrast, if there is some variable other than gender or class that differs across dormitories, sampling by floor may produce a more representative sample.

- 1.24 Row 16: 2323 6737 5131 8888 1718 0654 6832 4647 6510 4877 (a) Row 17: 4579 4269 2615 1308 2455 7830 5550 5852 5514 7182 Row 18: 0989 3205 0514 2256 8514 4642 7567 8896 2977 8822 Row 19: 5438 2745 9891 4991 4523 6847 9276 8646 1628 3554 Row 20: 9475 0899 2337 0892 0048 8033 6945 9826 9403 6858 Row 21: 7029 7341 3553 1403 3340 4205 0823 4144 1048 2949 Row 22: 8515 7479 5432 9792 6575 5760 0408 8112 2507 3742 Row 23: 1110 0023 4012 8607 4697 9664 4894 3928 7072 5815 Row 24: 3687 1507 7530 5925 7143 1738 1688 5625 8533 5041 Row 25: 2391 3483 5763 3081 6090 5169 0546 Note: All sequences above 5000 are discarded. There were no repeating sequences. 089 189 289 389 489 589 689 789 889 989 (b) 1089 1189 1289 1389 1489 1589 1689 1789 1889 1989 2089 2189 2289 2389 2489 2589 2689 2789 2889 2989 3089 3189 3289 3389 3489 3589 3689 3789 3889 3989 4089 4189 4289 4389 4489 4589 4689 4789 4889 4989
 - (c) With the single exception of invoice #0989, the invoices selected in the simple random sample are not the same as those selected in the systematic sample. It would be highly unlikely that a random process would select the same units as a systematic process.
- 1.25 (a) A stratified sample should be taken so that each of the four strata will be proportionately represented.
 - (b) Since the stratum may differ in the invoice amount, it may be more important to sample a larger percentage of invoices in stratum 1 and stratum 2, and smaller percentages in stratum 3 and stratum 4. For example, 50/5000 = 1% so 1% of 500 = 5 invoices should be selected from stratum 1; similarly 10% = 50 should be selected from stratum 2, 20% = 100 from stratum 3, and 69% = 345 from stratum 4.
 - (c) It is not simple random sampling because, unlike the simple random sampling, it ensures proportionate representation across the entire population.
- 1.26 Before accepting the results of a survey of college students, you might want to know, for example:

Who funded the survey? Why was it conducted? What was the population from which the sample was selected? What sampling design was used? What mode of response was used: a personal interview, a telephone interview, or a mail survey? Were interviewers trained? Were survey questions field-tested? What questions were asked? Were they clear, accurate, unbiased, valid? What operational definition of "vast majority" was used? What was the response rate? What was the sample size?

- 1.27 (a) Possible coverage error: Only employees in a specific division of the company were sampled.
 - (b) Possible nonresponse error: No attempt is made to contact nonrespondents to urge them to complete the evaluation of job satisfaction.
 - (c) Possible sampling error: The sample statistics obtained from the sample will not be equal to the parameters of interest in the population.
 - (d) Possible measurement error: Ambiguous wording in questions asked on the questionnaire.

42 Chapter 1: Defining and Collecting Data

- 1.28 The results are based on an online survey. If the frame is supposed to be small business owners, how is the population defined? This is a self-selecting sample of people who responded online, so there is an undefined nonresponse error. Sampling error cannot be determined since this is not a random sample.
- 1.29 Before accepting the results of the survey, you might want to know, for example: Who funded the study? Why was it conducted? What was the population from which the sample was selected? What was the frame being used? What sampling design was used? What mode of response was used: a personal interview, a telephone interview, or a mail survey? Were interviewers trained? Were survey questions field-tested? What other questions were asked? Were they clear, accurate, unbiased, and valid? What was the response rate? What was the margin of error? What was the sample size?
- 1.30 Before accepting the results of the survey, you might want to know, for example: Who funded the study? Why was it conducted? What was the population from which the sample was selected? What sampling design was used? What mode of response was used: a personal interview, a telephone interview, or a mail survey? Were interviewers trained? Were survey questions field-tested? What other questions were asked? Were the questions clear, accurate, unbiased, and valid? What was the response rate? What was the margin of error? What was the sample size? What frame was used?
- 1.31 A population contains all the items of interest whereas a sample contains only a portion of the items in the population.
- 1.32 Categorical random variables yield categorical responses such as yes or no answers. Numerical random variables yield numerical responses such as your height in inches.
- 1.33 Discrete random variables produce numerical responses that arise from a counting process. Continuous random variables produce numerical responses that arise from a measuring process.
- 1.34 Both nominal scaled and ordinal scaled variables are categorical variables but no ranking is implied in nominal scaled variable such as male or female while ranking is implied in ordinal scaled variable such as a student's grade of A, B, C, D and F.
- 1.35 Both interval scaled and ratio scaled variables are numerical variables in which the difference between measurements is meaningful but an interval scaled variable does not involve a true zero such as standardized exam scores while a ratio scaled variable involves a true zero such as height.
- 1.36 Microsoft Excel could be used to perform various statistical computations that were possible only with a slide-rule or hand-held calculator in the old days.
- 1.37 (a) The population of interest was all small business owners and/or employees responsible for information technology (IT).
 - (b) The sample was the 1,232 small businesses, having between 2 and 99 employees, drawn from e-Rewards's online business panel of companies.

Statistics for Managers Using MS Excel Global 7th Edition Levine Solutions Manual

Full Download: https://alibabadownload.com/product/statistics-for-managers-using-ms-excel-global-7th-edition-levine-solutions-managers-ms-excel-global-7th-edition-levine-solutions-managers-ms-excel-global-7th-edition-solutions-managers-ms-excel-global-7th-editions-managers-ms-excel-global-7th-editions-ms-excel-global-7th-editions-ms-excel-global-7th-editions-ms-ex

Solutions to End-of-Section and Chapter Review Problems 43

- 1.38 The answers to this question depend on which article and its corresponding data set is being selected.
- 1.39 (a) The population of interest was all adults living in the United States, aged 18 and older.
 - (b) The sample was the 1,016 adults, aged 18 and older, living in all 50 U.S. states and the District of Columbia that responded to the telephone interviews conducted on April 9–12, 2012.
- 1.40 The answers to this question depend on which data set is being selected.
- 1.41 (a) Categorical variable: Which of the following best describes this firm's primary business?
 (b) Numerical variable: On average, what percent of total monthly revenues are e-commerce revenues?
- 1.42 (a) The population of interest was the collection of all the 10,000 benefitted employees at the University of Utah when the study was conducted.
 - (b) The sample consisted of the 3,095 benefitted employees participated in the study.
 - (c) gender: categorical; age: numerical; education level: numerical; marital status: categorical; household income: numerical; employment category: categorical

1.43	(a)	(i)categorical	(iii) numerical, discrete
		(ii)categorical	(iv) categorical

- (b) The answers will vary.
- (c) The answers will vary.