Full Download: http://testbanklive.com/download/college-physics-volume-1-11th-edition-serway-solutions-manual/ Topic 1

Topic 1

Units, Trigonometry, and Vectors

QUICK QUIZZES

1.1Choice (c). The largest possible magnitude of the resultant occurs when the two vectors are in the same direction. In this case, the magnitude of the resultant is the sum of the magnitudes of \vec{A} and $\vec{B}: R = A + B = 20$ units. The smallest possible magnitude of the resultant occurs when the two vectors are in opposite directions, and the magnitude is the difference of the magnitudes of \vec{A} and $\vec{B}: R = |A - B| = 4$ units.

1	.2	

Vector	x-component	y-component
Ā	_	+
B	+	_
$\vec{A} + \vec{B}$	_	_

1.3Vector \vec{B} . The range of the inverse tangent function includes only the first

and fourth quadrants (i.e., angles in the range $-\pi/2 < \theta < \pi/2$). Only

© 2018 Cengage Learning. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

Full download all chapters instantly please go to Solutions Manual, Test Bank site: testbanklive.com

vector \mathbf{B} has an orientation in this range.

ANSWERS TO EVEN NUMBERED CONCEPTUAL QUESTIONS

- **1.2** Atomic clocks are based on the electromagnetic waves that atoms emit. Also, pulsars are highly regular astronomical clocks.
- **1.4** (a) ~0.5 lb \approx 0.25 kg or ~10⁻¹ kg
 - **(b)** $\sim 4 \text{ lb} \approx 0.25 \text{ kg or } \sim 10^{\circ} \text{ kg}$
 - (c) $\sim 4\ 000\ \text{lb} \approx 2\ 000\ \text{kg}\ \text{or}\ \sim 10^3\ \text{kg}$
- **1.6** Let us assume the atoms are solid spheres of diameter 10^{-10} m. Then, the volume of each atom is of the order of 10^{-30} m³. (More precisely, volume = $4\pi r^3/3$.) Therefore, since $1 \text{ cm}^3 = 10^{-6} \text{ m}^3$, the number of atoms in the 1 cm³ solid is on the order of $10^{-6}/10^{-30} = 10^{24}$ atoms. A more precise calculation would require knowledge of the density of the solid and the mass of each atom. However, our estimate agrees with the more precise calculation to within a factor of 10.
- **1.8** Choice (d). For an angle θ from 0° to 360°, the sine and cosine functions take the values

$$-1 \le \sin \theta \le 1$$
 and $-1 \le \cos \theta \le 1$

1.10 In the metric system, units differ by powers of ten, so it's very easy

and accurate to convert from one unit to another.

- 1.12 Both answers (d) and (e) could be physically meaningful. Answers (a),(b), and (c) must be meaningless since quantities can be added or subtracted only if they have the same dimensions.
- **1.14** The correct answer is (a). The second measurement is more precise but, given the number of reported significant figures, each measurement is consistent with the other.
- **1.16** The components of a vector will be equal in magnitude if the vector lies at a 45° angle with the two axes along which the components lie.

ANSWERS TO EVEN NUMBERED PROBLEMS

1.2	(a)	L / T ²	(b)	L			
1.4	All three equations are dimensionally incorrect.						
1.6	(a)	kg ∙ m/s			(b)	Ft = p	
1.8	58						
1.10	(a)	22.6			(b)	22.7	(c) 22.6 is more reliable
1.12	(a)	3.00×10^{-10}) ⁸ m/s		(b)	2.997 9 >	$\times 10^8 \text{ m/s}$
	(c)	2.997 92	5×10^{8}	m/s			
1.14	(a)	346 m ² ±	- 13 m ³		(b)	66.0 m ±	- 1.3 m

3

1.16	(a)	797	(b)	1.1	(c)	17.66				
1.18	132	m ²								
1.20	3.09	cm/s								
1.22	(a)	$5.60 \times 10^2 \text{ km} = 5.60 \times 10^2 \text{ km}$	$0^{5} m =$	5.60×10^{-10}	0^7 cm					
	(b)	(b) 0.491 2 km = 4.912×10^4 cm								
	(c)	(c) $6.192 \text{ km} = 6.192 \times 10^3 \text{ m} = 6.192 \times 10^5 \text{ cm}$								
	(d)	(d) 2.499 km = 2.499×10^3 m = 2.499×10^5 cm								
1.24	10.6	km/L								
1.26	9.2 nm/s									
1.28	$2.9 \times 10^2 \text{ m}^3 = 2.9 \times 10^8 \text{ cm}^3$									
1.30	$2.57 \times 10^{6} \text{ m}^{3}$									
1.32	$\sim 10^8$ steps									
1.34	~10 ⁸ people with colds on any given day									
1.36	(a)	$4.2 \times 10^{-18} \text{ m}^3$	(b)	~10 ⁻¹ m	3	(c)	$\sim 10^{16}$ cells			
1.38	10^{-14}	⁴ kg								
1.40	2.2 1	m								
1.42	8.1 0	cm								

1.44
$$\Delta s = \sqrt{r_1^2 + r_2^2 - 2r_1r_2\cos(\theta_1 - \theta_2)}$$

 $^{\odot}$ 2018 Cengage Learning. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

College Physics Volume 1 11th Edition Serway Solutions Manual

Full Download: http://testbanklive.com/download/college-physics-volume-1-11th-edition-serway-solutions-manual/ Topic 1 5

τορις τ			0			
1.46	2.33 m					
1.48	(a) 1.50 m	(b) 2.60 m				
1.50	8.60 m					
1.52	$1.44 \times 10^3 \text{ m}$					
1.54	(a) 6.1 units at $\theta = +113^{\circ}$	(b) 15 units at $\theta = +23^{\circ}$				
1.56	(a) 484 km	(b) 18.1° N of W				
	(c) Because of Earth's curvature, the plane does not follow straight					
	lines.					
1.58	$\vec{\mathbf{R}} = 9.5$ units at 57° above the	e + x-axis				
1.60	(a) 13.4 m (b) 19.9 m					
1.62	1.31 km northward, 2.81 km e	eastward				
1.64	(a) 10.0 m (b) 15.7 m	(c) 0				
1.66	42.7 yards					
1.68	788 mi at 48.0° N of E					
1.70	(a) 185 N at 77.8° from the <i>x</i> -	axis (b) 185 N at 258° from the <i>x</i> -ax	is			
1.72	(a) 1.609 km/h	(b) 88 km/h (c) 16 km	n/h			
1.74	(a) 7.14×10^{-2} gal/s	(b) $2.70 \times 10^{-4} \text{ m}^3/\text{s}$				
	(c) 1.03 h					

© 2018 Cengage Learning. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

Full download all chapters instantly please go to Solutions Manual, Test Bank site: testbanklive.com